arXiv:1108.1501v1 [math.DS] 6 Aug 2011

Beyond Periodicity for Piecewise Contracting Maps

E. Catsigeras[1], P. Guiraud[2], A. Meyroneinc[3] and E. Ugalde[4]

August 9, 2011

[1] Instituto de Matemática y Estadística Rafael Laguardia, Universidad de la República,
Montevideo, Uruguay
eleonora@fing.edu.uy

[2] Centro de Investigación y Modelamiento de Fenómenos Aleatorios Valparaíso, Universidad de Valparaíso,
Valparaíso, Chile
pierre.guiraud@uv.cl

[3] Departamento de Matemáticas, Instituto Venezolano de Investigaciones Científicas,
Apartado 20632, Caracas 1020A, Venezuela
ameyrone@ivic.gob.ve

[4] Instituto de Física, Universidad Autónoma de San Luis Potosí,
78000 San Luis Potosí, Mexico
ugalde@ifisica.uaslp.mx

Abstract

We study a wide class of dynamical systems defined by the iteration of piecewise contracting maps on compact and locally connected metric spaces. We consider non–isolated discontinuities and finitely many pieces of continuity in any finite or infinite dimensional space. We study the topological structure of the global attractors, non–wandering and limit sets associated to the orbits that do not intersect the frontiers of the different pieces of contraction. First, we obtain general results proving several new theorems and generalizing some old ones. We revisit the known strong hypothesis implying that the global attractor is made of a finite number of periodic orbits. We also give mild conditions on this class of maps that are sufficient for the total disconnection of the attractor and for its good recurrence properties. Second, we construct a series of examples for the mostly unknown case where the attractor intersects the frontiers. Thereby we show that the attractor may be finite, countably infinite or uncountable; it may include non–recurrent and also non–wandering points; and even when all the points in the attractor are non–wandering we prove that it does not necessarily coincide with the closure of all the admissible limit sets. This diversity of possible asymptotic behaviours is mostly interesting for models with milder structures than those commonly adopted.

Keywords: Piecewise contraction, Periodic points, Attractor, Recurrence.

MSC 2010: 37B25, 37B20, 54C08, 37N99

1 Introduction

The traditional concern of dynamical systems theory being mainly on sufficiently smooth flows and maps, the great majority of the results for piecewise smooth and piecewise continuous dynamical systems have been for long essentially restricted to low dimensional phenomena [19]. During the last two decades however, an increasing attention has been paid to higher dimensions. Particularly noticeable is the case of weakly coupled piecewise expanding maps for which new techniques to study ergodic properties in infinite dimensional phase spaces have been developed, see [22] and references therein. We can also mention the case of piecewise isometries for which zero topological entropy has been proved in any dimension [5], despite non–trivial recurrence properties [18, 24]. Applied fields are also a motivation for such new developments, piecewise continuous dynamical systems of arbitrary dimension being common in the modelling of systems in the natural and social sciences. To give a few examples only, they are used in control engineering as models for electro–mechanical systems [2, 14, 26], in biology as models for neural networks [7, 9, 10, 25] and for genetic regulatory networks [6, 11, 13, 23], and in social sciences as models in economics [28] and management [26]. In particular, within the wide class of piecewise continuous systems, piecewise contractions in any dimension appear in models where dissipative effects coexist with threshold–dependent interactions. We can mention discrete time models [10, 11, 23] and Poincaré return maps of continuous time models [7, 9, 13, 25] of neuronal and genetic networks. Despite the interest of piecewise contracting maps in physics and biology, less attention has been paid in mathematics to piecewise contracting maps than to their expanding and isometric counterparts.

For some very specific families of piecewise contracting maps it has been established that generically the asymptotic dynamics is periodic. It is the case for one–dimensional injective maps [3], two–dimensional affine maps [4], arbitrary dimensional affine maps with convex continuity pieces [10, 12, 23] and more general real maps satisfying a so–called separation property [7, 8]. All the proofs of periodicity need the limit set of the dynamics to be at positive distance of the discontinuity set, a property that indeed turns to be generic in those special cases, but is not proved in more general contexts. Also, efforts being concentrated on the generic situation, very little is known when the limit set intersects the discontinuity set. The few examples considering this situation show an asymptotic dynamics which is an irrational rotation on a Cantor set [11, 17, 21]. But in general the questions of the possible topologies of the asymptotic sets and the dynamics on them when they intersect the discontinuity set remain open. The study of these questions is a preliminary step in the analysis of bifurcations [1, 11] and the investigation of quantitative properties such as complexity [23], chaos in the sense of [15, 16], and topological entropy [29] in piecewise contracting systems.

In this paper we investigate the limits of a general theory of the dynamics of piecewise contracting maps. We start with some general results concerning the topological and recurrence properties of the attractor of any piecewise contracting map defined on a compact and locally connected metric space. Because of the discontinuous character of the dynamics, we need first to revise the concept of attractor as well as those of limit set and non–wandering set. After providing general results about the recurrence properties of the attractor, we prove, in a context that encompasses the models considered so far, the asymptotic periodicity when the attractor is at positive distance of the discontinuity set. We do not pretend to prove here that the latter condition is generic. Admitting this condition though, the proof of periodicity is not direct since we cannot repeat usual arguments of connectedness that are true in the particular situations where genericity is proved. From the point of view of applications, inspired by [10], we propose to answer the question of the asymptotic periodicity of almost any kind of model by verifying that the attractor is at positive distance of the discontinuity set instead of verifying that the model belongs to a particular class of piecewise contracting maps and that, moreover, it is typical in that class.

As mentioned above, for the piecewise contracting systems considered so far in the literature, it has been observed that the long term dynamics is concentrated on a finite union of transitive components that are either periodic orbits or Cantor sets. After giving some sufficient conditions for the attractor being totally disconnected, by mean of (counter–)examples we prove that this scenario is in fact too restricted. Indeed, we show that – when the attractor intersects the discontinuity set – the transitive components of the attractor can be finite but not periodic[1], infinite but countable, uncountable but not restricted to a Cantor set. For instance, we show for the latter case that the attractor can consist of a unique transitive component which is neither a Cantor set nor a periodic orbit, but the union of a Cantor set with a countably infinite set. On the other hand, some of our examples illustrate diverse unexpected phenomena concerning the recurrence properties of the attractor, as they can be observed in some continuous dynamical systems having expanding directions near or along the attractor, such as the presence of wandering points. Altogether, our examples give insight on the variety of possible attractors for piecewise contracting maps and, in particular, they obsolete the extrapolation of the observed dichotomy according to which the asymptotic dynamics is either supported by a finite set of periodic orbits or by an uncountable set which is essentially a Cantor set.

The paper is organized as follows. In Section 2, we define the limit set, the non–wandering set and the attractor for general piecewise contracting maps. We then study the relations between these sets and prove the general theorems about the topology of the attractor. In Section 3, we present and study the series of examples illustrating the diverse attractors and phenomena that piecewise contracting dynamics may exhibit when their attractor intersects the discontinuity set. Finally, in Section 4 we present some final comments and open questions which may contribute to define new lines of research on this subject.

---

[1] When the attractor intersects the discontinuity set, according to the definition of the map on the latter, the attractor can fail to be invariant (and thus to be periodic) while being the ω–limit set of some point (and therefore being transitive).

2 Piecewise contracting maps

2.1 Definitions

Definition 2.1. Let X be a compact and locally connected space equipped with a distance d. We say that the map $f : X \mapsto X$ is *piecewise contracting* if there exists a finite collection of $N \geqslant 2$ non–empty open subsets $\{X_i\}_{i=1}^N$, such that:

1) $X = \bigcup_{i=1}^N \overline{X}_i$ and $X_i \cap X_j = \emptyset$ for all $i \neq j \in \{1, \dots, N\}$,

2) There exists $\lambda \in (0,1)$, called *a contracting rate,* such that for every $i \in \{1, \dots, N\}$,

$$d(f(x), f(y)) \leqslant \lambda d(x,y) \qquad \forall\, x, y \in X_i. \tag{1}$$

3) The set $\tilde{X} := \bigcap_{n=0}^{+\infty} f^{-n}(X \setminus \Delta)$, where $\Delta := X \setminus (\bigcup_{i=1}^N X_i)$, is non empty.

Note that a piecewise contracting map is Lipchitz continuous in each of the open sets X_i. All the possible discontinuities are contained in the set Δ, which is actually a meager set equal to the union of the borders of the X_i. In concrete examples we may encounter any kind of definition for f on the set Δ. In order to obtain results shared by all the piecewise contracting maps (i.e. independent of the definition of f on Δ), we restrict our set of initial conditions to the set $\tilde{X}$ of points whose orbits do not intersect Δ. This is why we consider the condition 3) which is usually a soft restriction, since the set $\tilde{X}$ is dense in X under mild hypothesis on the map f (see Theorem 2.8).

In what follows we will focus on the topological properties of the sets supporting the asymptotic dynamics, namely the limit set, the non–wandering set and the attractor. Because of the discontinuous nature of the dynamics, we have to adapt to the present context the standard definitions of these notions.

Definition 2.2. Let $f : X \to X$ be a piecewise contracting map. We say that $y \in X$ is an *ω–limit point* of $x \in \tilde{X}$, if there exists a divergent sequence $\{n_k\}_{k\in\mathbb{N}}$ such that $\lim\limits_{k\to\infty} f^{n_k}(x) = y$. We call *$\omega$–limit set* and denote $\omega(x)$ the set of all the ω–limit points of x. We call *limit set* of f the set

$$L := \overline{\bigcup_{x\in\tilde{X}} \omega(x)}.$$

Due to the compactness of the space, $\omega(x) \neq \emptyset$ for all $x \in X$, and in particular for all $x \in \tilde{X}$. Therefore, as $\tilde{X} \neq \emptyset$, the limit set $L \neq \emptyset$. As in the standard definition of recurrence for continuous maps, a point $x \in \tilde{X}$ is called *recurrent* if $x \in \omega(x)$. Trivially, the set $\mathcal{R}$ of all the recurrent points in $\tilde{X}$, is contained in the limit set L. Nevertheless, the limit set can contain points in $\tilde{X}$ that are no recurrent, as shown in Example 3.2 of Section 3.

Definition 2.3. Let $f : X \to X$ be a piecewise contracting map. We say that a point $x \in X$ is *non–wandering* if for all $\epsilon > 0$ there exists a divergent sequence $\{n_k\}_{k\in\mathbb{N}}$ such that the ball $B(x,\epsilon)$ of center x and radius ϵ satisfies:

$$f^{n_k}(B(x,\epsilon) \cap \tilde{X}) \bigcap B(x,\epsilon) \neq \emptyset \qquad \forall\, k \in \mathbb{N}. \tag{2}$$

We denote Ω the *non–wandering set*, defined as the set of all the non wandering points. We say that a point is *wandering* if it does not belong to Ω.

The principal difference with the standard definition of wandering point for continuous maps is in that the initial conditions returning to $B(x,\epsilon)$ must belong to $\tilde{X}$, as we disregard all the orbits that intersect Δ (but x can be a point of Δ). If we would not, the arbitrary dynamical behaviour of those initial conditions in

$X \setminus \tilde{X}$, after visiting Δ, may change the non–errancy quality of the other points of the space. By this way, we ensure that our definition of the non–wandering set is independent of all the possible arbitrary choices of f on Δ. It is standard to check that Ω is compact and contains L (see Theorem 2.9), and thus $\Omega \neq \emptyset$.

As for some other bifurcating continuous dynamical systems, the long term global dynamics for piecewise contracting maps may takes place on a larger set (possibly of different topological type) than the limit set L, and even than the non–wandering set Ω (we give concrete examples of these phenomena in Section 3). This larger set Λ, which we call attractor, is defined as follows:

Definition 2.4. Let $f : X \to X$ be a piecewise contracting map and consider the family of transformations $\{F_i\}_{i\in\{1,\dots,N\}}$ defined for all $A \subset X$ by $F_i(A) = \overline{f(A \cap X_i)}$. Let $n \geqslant 1$. We say that A is an *atom of generation* n if

$$A = F_{i_n} \circ F_{i_{n-1}} \circ \cdots \circ F_{i_1}(X) \tag{3}$$

for some $i_1, i_2, \dots, i_n \in \{1, \dots, N\}$. Denote $\mathcal{A}_n$ the set of all the atoms of generation n, and let

$$\Lambda_n := \bigcup_{A \in \mathcal{A}_n} A \qquad \forall\, n \geqslant 1.$$

We call *attractor* of (X, f) to the set

$$\Lambda := \bigcap_{n \geqslant 1} \Lambda_n.$$

For a dynamical system defined by iterations of a continuous transformation $g : X \to X$, it is usually called attractor a forward invariant and compact subset Λ such that $\Lambda = \bigcap_{n\geqslant 0} g^n(V)$, where V is an invariant neighborhood of Λ. For piecewise contractive maps, we need to introduce the above concept of atom (that appears also in [7, 8, 9] and [23] in a different form) in order to disregard the asymptotic sets resulting from the arbitrary dynamics in Δ. This concept allows to generalize the notion of attractor to our case (as explained bellow) and guaranties its computation is independent of the definition of f in Δ, although it could intersect Δ.

A property we will often refer to is that any atom is included in an atom of the previous generation. It implies the inclusion $\Lambda_{n+1} \subset \Lambda_n$ for all $n \geqslant 1$. Since we suppose $\tilde{X}$ non–empty, there exists always a non–empty atom in $\mathcal{A}_n$. The sets Λ_n, and therefore Λ, are thus non–empty compact sets. Besides, if for some $n_0 \geqslant 1$ the set Λ_{n_0} is included in $\tilde{X}$ where all the iterates of f are continuous, then $\Lambda_{n+1} = f(\Lambda_n)$ for all $n \geqslant n_0$. It follows that the compact set Λ of Definition 2.4, is an attractor also in the sense of the standard definition for continuous systems, because $\Lambda = \bigcap_{n\geqslant 0} f^n(\Lambda_{n_0})$, where Λ_{n_0} is a invariant neighborhood of Λ.

The above arguments show that Definition 2.4 is a generalization to the case of piecewise contracting maps (and more generally to piecewise continuous maps), of the notion of attractor of a continuous system. However, since in general Λ intersects the set Δ, depending on the definition of f in Δ, the attractor (as well as the limit set and the non wandering set) could fail to be invariant, as we will see in Section 3. On the other hand, from the definition of atoms it follows $f(\Lambda_n \setminus \Delta) \subset \Lambda_{n+1} \subset \Lambda_n$, which implies $f(\Lambda \setminus \Delta) \subset \Lambda$. In other words, the image of a point of the attractor which is not in Δ belongs to the attractor. We say that the attractor is *essentially invariant*. Nevertheless, when the attractor intersects Δ it may be impossible to define a sub–system $f|_\Lambda : \Lambda \mapsto \Lambda$ and that is why we adopt now the following definition of transitivity.

Definition 2.5. We say that a compact set $Y \subset X$ is *transitive* if there exists $x \in \tilde{X}$ such that $Y = \omega(x)$.

When the attractor is not transitive, it may nevertheless contain transitive sets that we call *transitive components*. Note that all transitive sets are contained in the limit set L.

We close this subsection, with a remark and some technical definitions about the atoms that we will refer to in the sequel of the paper.

Remark 2.6. The atoms are not necessarily connected sets. Even if the metric space (X, d) is for instance an Euclidean compact space of dimension two, the intersection $A \cap X_i$ may be disconnected, regardless whether A and X_i are both connected sets. In particular, we found piecewise contractive maps defined in a compact square of $\mathbb{R}^2$, that are affine in each contracting piece, and that exhibit atoms of arbitrarily large generations with infinitely many connected components (see Appendix 5). However, it is easy to check that the diameter of the atoms converges to zero when their generation tends to infinity. The local connectedness of the space X ensures then that any atom of sufficiently large generation is contained in a connected ball, even if the atom itself is not necessarily a connected set. This observation will help the proof of our Theorems 2.9, 2.10 and 2.12 in the next subsection.

Definition 2.7. 1) The word of indexes $(i_1, i_2, \dots, i_n)$ with length n, defining each atom $A \in \mathcal{A}_n$ according to Equality (3), is called the *n–itinerary* of A. We say that *two atoms are differently indexed* if their respective n-itineraries are different, even if they coincide as subsets of X, and thus, are the same atom.

2) We say that a map is *atomically one–to–one* if two differently indexed atoms of the same generation, either coincide or are disjoint.

3) We say that a word of indexes $(i_1, i_2, \dots, i_n)$ is *admissible,* if it is the n-itinerary of a non-empty atom of $\mathcal{A}_n$.

2.2 General properties

We first prove that under mild hypothesis on the piecewise contracting map f, the set $\tilde{X}$ is dense in X. The proof works also for piecewise continuous maps but we enunciate the theorem in our context.

Theorem 2.8. *Let $f : X \to X$ be a piecewise contracting map such that for all $i \in \{1, 2, \dots, N\}$ the restricted map $f|_{X_i} : X_i \mapsto f(X_i)$ is an homeomorphism. Then $\tilde{X} := \bigcap_{n=0}^{+\infty} f^{-n}(X \setminus \Delta)$ is dense in X.*

Proof. The metric space X being compact it is complete, and therefore it is a Baire space. Since

$$\tilde{X} = \bigcap_{k=0}^{+\infty} \tilde{X}_k \qquad \text{where} \quad \tilde{X}_k := \bigcap_{n=0}^{k} f^{-n}(X \setminus \Delta) \quad \forall\, k \in \mathbb{N},$$

to prove the theorem it is enough to show that the $\tilde{X}_k$ are open and dense in X, as by definition all countable intersection of open dense sets is dense. It is easy to show the following relation:

$$\tilde{X}_{k+1} = \bigcup_{i=1}^{N} (f|_{X_i})^{-1}(\tilde{X}_k) \quad \forall\, k \in \mathbb{N}, \tag{4}$$

noting that $\bigcup_{i=1}^{N} (f|_{X_i})^{-1}(\tilde{X}_k) = \{x \in X \setminus \Delta \ : \ f(x) \in \tilde{X}_k\}$ and using the definition of the $\tilde{X}_k$. We use it to show by induction that $\tilde{X}_k$ is open and dense in X for all $k \in \mathbb{N}$. As $\tilde{X}_0 = \bigcup_{i=1}^{N} X_i = \text{int}(X)$ it is true for $k = 0$. Now suppose that $\tilde{X}_{k_0}$ is open and dense in X for a $k_0 \in \mathbb{N}$ and let $i \in \{1, \dots, N\}$. First, $(f|_{X_i})^{-1}(\tilde{X}_{k_0}) = (f|_{X_i})^{-1}(\tilde{X}_{k_0} \cap f(X_i))$ is open. Indeed, since $f|_{X_i}$ is an homeomorphism, $\tilde{X}_{k_0} \cap f(X_i)$ is the intersection of two open sets whose pre–image is open. Second, as $f(X_i)$ is open and $\tilde{X}_{k_0}$ is dense in X, we have also that $\tilde{X}_{k_0} \cap f(X_i)$ is dense in $f(X_i)$. It results that $(f|_{X_i})^{-1}(\tilde{X}_{k_0})$ is dense in X_i. It follows then from (4) that $\tilde{X}_{k_0+1}$ is open, and it is dense since

$$\overline{\tilde{X}_{k_0+1}} = \bigcup_{i=1}^{N} \overline{(f|_{X_i})^{-1}(\tilde{X}_{k_0})} = \bigcup_{i=1}^{N} \overline{X_i} = X.$$

□

It is well known that any continuous dynamical system (X, g) satisfies $\Lambda \supset \Omega \supset L$, where $\Lambda := \bigcap_{n \in \mathbb{N}} g^n(X)$ is the attractor, Ω is the non–wandering set and $L := \overline{\bigcup_{x \in X} \omega(x)}$ is the limit set. The following Theorem 2.9 states, in particular, that it is also true for piecewise contracting maps.

Theorem 2.9. *Let $f : X \to X$ be a piecewise contracting map. Denote L its limit set, Ω its non–wandering set and Λ its global attractor. Then,*

i) $L \subset \Omega \subset \Lambda$.[2]

ii) Any non–wandering point in the interior of $\tilde{X}$ is recurrent.

---

[2] It is actually true for piecewise continuous maps in general, since the proof requires neither the existence of a contraction rate nor any kind of connectedness of X.

For any continuous system on a compact metric space, provided it is uniformly contracting, the three sets describing the asymptotic dynamics, coincide: $\Lambda = \Omega = L$. This result follows directly from the Banach fixed point Theorem. In other words, to have $\Lambda \neq \Omega$ or $\Omega \neq L$ for a continuous system evolving in a compact metric space, it is necessary some topological expanding property, even if it is partial, or just the loss of an uniform contracting rate. For instance, for some bifurcating diffeomorphisms acting on a compact two–dimensional manifold, and exhibiting an homoclinic tangency of a dissipative saddle periodic point, the limit set and the non–wandering set are not equal [27]. This last phenomenon also appears for piecewise contractive systems, but without the presence of any expanding property, and without the loss of the uniform contracting rate $\lambda \in (0, 1)$. In fact, in the examples 3.2 and 3.3 of Section 3, we show the existence of piecewise contractive maps such that $\Lambda \neq \Omega$ or $\Omega \neq L$. In this sense, the discontinuities of a piecewise contracting map act as the generators of a topological *virtual* expansion, even if the map is uniformly contracting in all its continuity pieces.

As a consequence of ii), a point in the attractor which is in the interior of $\tilde{X}$ and is not recurrent, is wandering. Thus, in the particular case where $\Lambda \setminus \mathcal{R}$ is non–empty and is a subset of the interior of $\tilde{X}$, all the non–recurrent points of Λ are wandering, which implies together with i) that $\Omega \setminus \mathcal{R} = \emptyset$ and $\mathcal{R} = \Omega = L$. Now, Λ could not contain recurrent points, or Λ could intersect Δ, so the condition $\Lambda \setminus \mathcal{R} \subset \text{int}(\tilde{X})$ could be quite strong or not be satisfied (recall that recurrent points are in $\tilde{X}$). Nevertheless, we can relax this condition to obtain similar results. Indeed, if $\Lambda \setminus L$ is non–empty and is a subset of $\text{int}(\tilde{X})$, then $\Omega \setminus L$ is also a subset of $\text{int}(\tilde{X})$ and is empty, since if not, from the property ii), we should have $\Omega \setminus L \subset \mathcal{R}$, which is a contradiction. It results that under the condition $\Lambda \setminus L \subset \text{int}(\tilde{X})$, we have $\Omega = L$ and $\Lambda \setminus L$ contains only wandering points. This is virtually the same result, except that we do not know if the limit points are all recurrent.

In Section 3, we construct Example 3.3 for which $\Lambda \setminus L$ has infinitely many points, all of them in the interior of $\tilde{X}$. It proves the existence of piecewise contracting maps whose attractor has infinitely many wandering points. On the other hand, we also show that the hypothesis $\Lambda \setminus L \subset \text{int}(\tilde{X})$ is not redundant to obtain the thesis $L = \Omega$. This is done with Example 3.4 where we construct a piecewise contracting map which does not satisfies that hypothesis. In this example, $\Lambda \setminus L$ is infinite, all its points are non wandering, and $\Omega \neq L$.

Proof. of Theorem 2.9: First we prove that $L \subset \Omega$. Let $y \in L$. According to Definition 2.2, there exists $x \in \tilde{X}$ and a diverging sequence $\{n_k\}_{k\in\mathbb{N}}$ such that $\lim\limits_{k\to\infty} f^{n_k}(x) = y$. Thus, if we fix $\epsilon > 0$, there exists $k_0 \in \mathbb{N}$ such that $f^{n_{k_0+p}}(x)$ belongs to the ball $B(y, \epsilon)$ of center y and radius ϵ for all $p \in \mathbb{N}$. In other words, $f^{m_p}(f^{n_{k_0}}(x)) \in B(y, \epsilon)$ for all $p \in \mathbb{N}$, if $\{m_p\}_{p\in\mathbb{N}}$ is the diverging sequence defined by $m_p = n_{k_0+p} - n_{k_0}$. Since $f^{n_{k_0}}(x) \in B(y, \epsilon) \cap \tilde{X}$, it follows that $f^{m_p}(B(y, \epsilon) \cap \tilde{X}) \cap B(y, \epsilon) \neq \emptyset$ for all $p \in \mathbb{N}$. According to Definition 2.3 the point $y \in \Omega$, and we have proved that $L \subset \Omega$.

Now we prove that $\Omega \subset \Lambda$. Let $y \in \Omega$. By definition, for any fixed $\epsilon > 0$ there exists a diverging sequence $\{n_k(\epsilon)\}_{k\in\mathbb{N}}$ such that,

$$f^{n_k(\epsilon)}(B(y, \epsilon) \cap \tilde{X}) \cap B(y, \epsilon) \neq \emptyset \qquad \forall\, k \in \mathbb{N}.$$

Thus, we can construct a sequence of points $\{y_j\}_{j\in\mathbb{N}}$ and a diverging sequence of natural numbers $\{n_j\}_{j\in\mathbb{N}}$ such that

$$y_j \in f^{n_j}(B(y, 1/j) \cap \tilde{X}) \cap B(y, 1/j) \qquad \forall\, j \in \mathbb{N}.$$

On one hand, for any $j \in \mathbb{N}$ we have $y_j \in \Lambda_{n_j}$ because y_j belongs to $f^{n_j}(\tilde{X})$ which is contained in Λ_{n_j}. On the other hand, as $y_j \in B(y, 1/j)$ for all $j \in \mathbb{N}$, we have $\lim\limits_{j\to\infty} y_j = y$. Therefore, $\lim\limits_{j\to\infty} d(\Lambda_{n_j}, y) = 0$. As $\Lambda_{n+1} \subset \Lambda_n$ for all $n \in \mathbb{N}$, it follows that $\lim\limits_{n\to\infty} d(\Lambda_n, y) = 0$. We deduce that $d(\Lambda, y) = 0$, and $y \in \Lambda$ by compactness of Λ. This finish the proof of the statement i) of Theorem 2.9.

Let us prove now the statement ii). Let $y \in \text{int}(\tilde{X})$ be non wandering. Since $y \in \text{int}(\tilde{X})$, there exists $\epsilon_0 > 0$ such that for all $\epsilon \in (0, \epsilon_0]$ the ball $B(y, \epsilon)$ of center y and radius ϵ satisfies $B(y, \epsilon) \subset \tilde{X}$. Therefore, by definition of $\tilde{X}$,

$$f^n(B(y, \epsilon)) \subset \bigcup_{i=1}^{N} X_i \qquad \forall\, n \in \mathbb{N}. \tag{5}$$

Besides, since the space X is locally connected, we can choose ϵ_0 such that for all $\epsilon \in (0, \epsilon_0]$ the ball $B(y, \epsilon)$ is a connected set. So the image of the ball by any continuous map is also a connected set. Since f is continuous in $\bigcup_{i=1}^N X_i$ it follows from the inclusion (5) that $f^n(B(y, \epsilon))$ is a connected set for all $n \in \mathbb{N}$. On the other hand, from the same inclusion we deduce that $f^n(B(y, \epsilon))$ can be written as:

$$f^n(B(y, \epsilon)) = \bigcup_{i=1}^{N} (X_i \cap f^n(B(y, \epsilon)) \qquad \forall\, n \in \mathbb{N}.$$

But as the $X_i \cap f^n(B(y, \epsilon)$ are disjoint and open in $f^n(B(y, \epsilon)$ (recall that the X_i are disjoint and open), the connectedness of $f^n(B(y, \epsilon))$ implies the existence of $i_n \in \{1, \dots, N\}$ such that

$$\bigcup_{\substack{i=1 \\ i \neq i_n}}^{N} (X_i \cap f^n(B(y, \epsilon)) = \emptyset.$$

Thus we have shown that for each $n \in \mathbb{N}$, the set $f^n(B(y, \epsilon))$ is contained in a single continuity piece X_{i_n} of f. Therefore, for all $\epsilon \in (0, \epsilon_0]$ and for all $n \in \mathbb{N}$, the iterated map $f^n|_{B(y,\epsilon)}$ is contracting with contraction rate λ^n. Thus,

$$f^n(B(y, \epsilon)) \subset B(f^n(y), \lambda^n \epsilon) \qquad \forall\, n \in \mathbb{N}, \quad \forall\, \epsilon \in (0, \epsilon_0]. \tag{6}$$

Now, let $\epsilon \in (0, \epsilon_0]$. As y is non wandering and $B(y, \epsilon) \subset \tilde{X}$, there exists a diverging sequence $\{n_k\}_{k \in \mathbb{N}}$ such that

$$f^{n_k}(B(y, \epsilon/2)) \cap B(y, \epsilon/2) \neq \emptyset \qquad \forall\, k \in \mathbb{N}.$$

Then, from the inclusion (6) we obtain

$$B(f^{n_k}(y), \lambda^{n_k} \epsilon/2) \cap B(y, \epsilon/2) \neq \emptyset \qquad \forall\, k \in \mathbb{N}.$$

This implies that $d(f^{n_k}(y), y) \leqslant (\lambda^{n_k} + 1)\epsilon/2 < \epsilon$ for all $k \in \mathbb{N}$. Since ϵ was chosen arbitrary in $(0, \epsilon_0]$, we have proved that for all $\epsilon \in (0, \epsilon_0]$ there exists a diverging sequence $\{n_k\}_{k \in \mathbb{N}}$ such that $f^{n_k}(y) \in B(y, \epsilon)$ for all $k \in \mathbb{N}$, i.e y is recurrent. □

In the forthcoming Theorem 2.10, we show that the hypothesis $d(\Lambda, \Delta) \neq 0$ implies that all the asymptotic sets coincide and consist of a finite number of periodic points. It has been shown this hypothesis is typical in several context, since it is satisfied for Lebesgue almost all piecewise contracting affine maps (with the Lebesgue measure in the space of a finite number of real parameters) [4, 10, 12, 23] and C^0–topologically generic for maps satisfying a separation property[3] [3, 7, 8]. In all these works, the phase space is homeomorphic to a connected compact rectangle embedded in $\mathbb{R}^n$. Here we do not prove the condition $d(\Lambda, \Delta) \neq 0$ is generic, we rather admit this condition and give a proof of periodicity in a very general context.

Theorem 2.10. *Let $f : X \to X$ be a piecewise contracting map with attractor Λ. If $\Delta = \emptyset$ or if $d(\Lambda, \Delta) \neq 0$, then Λ is a finite union of periodic orbits.*

Note that together with Theorem 2.9, this theorem implies that each asymptotic set Λ, Ω and L is equal to $\mathcal{R}$. To help the proof, we first prove Lemma 2.11, which states that each atom of sufficiently large generation is included in a single continuity piece of f. As observed in Remark 2.6, in general the atoms may be disconnected sets, even more, then can have an infinite number of connected components. So, we can not simply argue that the atoms are necessarily contained in a single continuity piece, provided that they do not intersect Δ. Nevertheless, that assertion is certainly true under strong additional hypothesis, for instance if all the continuity pieces are convex and embedded in $\mathbb{R}^n$ or if the map is an affinity in each piece (as in [4, 10, 12, 23]), since in this case all the atoms of all generations are indeed connected.

Lemma 2.11. *If $\Delta = \emptyset$ or if $d(\Lambda, \Delta) \neq 0$, then there exists $n_0 \geqslant 1$ such that for any atom A of generation $n \geqslant n_0$ there is a $i \in \{1, \dots, N\}$ such that $A \subset X_i$.*

---

[3] The separation property means that the map is atomically one–to–one, see Definition 2.7.

Proof. Since any atom is included is an atom of the previous generation, it is enough to show that there exists $n_0 \geqslant 1$ such that the lemma is true for any atom of generation n_0.

Suppose $\Delta = \emptyset$ and let $\delta > 0$ be sufficiently small for any ball of X with radius smaller than δ being connected (recall that the space X is locally connected). Choose $n_0 \in \mathbb{N}$ such that the diameter of any atom of $\mathcal{A}_{n_0}$ is smaller than δ (recall that the diameter of the atoms converges to 0 when their generation goes to infinity). Let $A \in \mathcal{A}_{n_0}$ and B_δ a ball of radius δ that contains A. Since $\Delta = \emptyset$, the X_i form a partition of X and $B_\delta = \bigcup_{i=1}^N (B_\delta \cap X_i)$. As B_δ is a connected set and the X_i are open, by definition of connectedness, B_δ (and therefore A) cannot intersects two of these sets. This closes the proof in the case $\Delta = \emptyset$.

Suppose $\Delta \neq \emptyset$. We first show that under the hypothesis $d(\Lambda, \Delta) \neq 0$ there exists $n_1 \in \mathbb{N}$ such that $d(\Lambda_{n_1}, \Delta) \neq 0$. By contradiction suppose $d(\Lambda_n, \Delta) = 0$ for all $n \in \mathbb{N}$. Then by compactness of Λ_n and Δ their intersection is non empty, and then there exists a sequence $\{x_n\}_{n\in\mathbb{N}}$ such that $x_n \in \Lambda_n \cap \Delta$ for all $n \in \mathbb{N}$. On one hand, again by compactness of Δ, this sequence converges to $x \in \Delta$. On the other hand, using the inclusion $\Lambda_{n+1} \subset \Lambda_n$ and the compactness of these sets, we obtain $x \in \Lambda_n$ for all $n \in \mathbb{N}$, i.e $x \in \Lambda$ (for any $p \in \mathbb{N}$, the point x is the limit of the sequence $\{y_n^p\}_{n\in\mathbb{N}}$ of points of Λ_p defined by $y_n^p = x_{n+p}$ for all $n \in \mathbb{N}$). So we have $x \in \Delta \cap \Lambda$, which contradicts $d(\Lambda, \Delta) \neq 0$.

Let $\delta > 0$ be such that $d(\Lambda_{n_1}, \Delta) > \delta$ and sufficiently small for any ball of X with radius smaller than δ being connected. Actually, by the inclusion $\Lambda_{n+1} \subset \Lambda_n$, we have $d(\Lambda_n, \Delta) > \delta$ for all $n \geqslant n_1$, and as Λ_n is the union of all the atoms of generation n we deduce

$$d(A, \Delta) > \delta \qquad \forall\, A \in \mathcal{A}_n \quad \forall\, n \geqslant n_1. \tag{7}$$

Choose $n_0 \geqslant n_1$ such that the diameter of any atom of $\mathcal{A}_{n_0}$ is smaller than δ. Let $A \in \mathcal{A}_{n_0}$ and $x \in A$. Then, from Inequality (7) we obtain:

$$d(x, \Delta) > \delta \tag{8}$$

Therefore $x \in X \setminus \Delta$ and there exists $i \in \{1, \ldots, N\}$ such that $x \in X_i$. Now if we suppose by contradiction that A intersects $X \setminus X_i$, the connected ball $B(x, \delta)$ intersects ∂X_i, i.e there exists $z \in B(x, \delta) \cap \partial X_i \subset \Delta$. This contradicts Inequality (8), since $z \in \Delta$ and $d(x, z) < \delta$. □

Proof. of Theorem 2.10: **Step 1 (Preliminary results).** Let $n \geqslant n_0$, where n_0 is such that Lemma 2.11 is satisfied. Let $A \in \mathcal{A}_n$, then there exists $i \in \{1, \ldots, N\}$ such that $A \subset X_i$. Together with the compactness of A and the continuity of f in X_i, it implies that $f(A) = \overline{f(A \cap X_i)}$. Therefore, for all $n \geqslant n_0$ and $A \in \mathcal{A}_n$, the set $f(A)$ is an atom of generation $n+1$ and reciprocally, according to the definition of atom, any atom of generation $n+1$ is the image of an atom of generation n. It follows that

$$f(\Lambda_n) := \bigcup_{A \in \mathcal{A}_n} f(A) = \bigcup_{A \in \mathcal{A}_{n+1}} A = \Lambda_{n+1} \qquad \forall\, n \geqslant n_0. \tag{9}$$

Moreover, since any atom of generation $n+1$ is contained in an atom of generation n, we have that for all $A \in \mathcal{A}_{n+1}$ with $n \geqslant n_0$ there exists $A' \in \mathcal{A}_n$ such that $f(A) \subset A'$.

Step 2. From Step 1 we deduce that $f : \Lambda_{n_0} \to \Lambda_{n_0}$ induces a transformation $G : \mathcal{A}_{n_0} \to \mathcal{A}_{n_0}$ given by $G(A) = A'$ if $f(A) \subset A'$. All the orbits of this transformation are eventually periodic, because the collection $\mathcal{A}_{n_0}$ where it acts, is finite. Thus, the atoms of generation n_0 are classified in those that are periodic by the transformation F, and those that are not periodic but, anyway, are eventually periodic. We denote $\mathcal{P}_{n_0}$ to the finite and not empty sub-collection of atoms in $\mathcal{A}_{n_0}$ that are periodic under the transformation G. Let $p_0 \geqslant 1$ be a multiple of the periods of all the atoms of $\mathcal{P}_{n_0}$, and let $m_0 \geqslant 1$ be the number of different atoms in the family $\mathcal{A}_{n_0}$. Then we have:

$$G^{m_0}(A) \in \mathcal{P}_{n_0} \qquad \text{and} \qquad G^{p_0}(F^{m_0}(A)) = G^{m_0}(A) \quad \forall\, A \in \mathcal{A}_{n_0}.$$

From the definition of G, we deduce that $f^{p_0}(f^{m_0}(A)) \subset f^{m_0}(A)$ for all $A \in \mathcal{A}_{n_0}$, or equivalently (by Step 1)

$$f^{p_0}(A) \subset A \quad \forall\, A \in \mathcal{A}_{n_0+m_0}. \tag{10}$$

From Lemma 2.11, it follows that f^{p_0} is contracting in any atom of generation greater that n_0. The atoms being compact, using the Banach fixed point theorem, we deduce from (10) that f admits a unique periodic point in each atom of $\mathcal{A}_{n_0+m_0}$.

Step 3. Now we show that all the points of Λ are periodic. Recalling that $\Lambda_{n+1} \subset \Lambda_n$ for any $n \geqslant 1$, we can write

$$\Lambda = \bigcap_{n\in\mathbb{N}} \Lambda_n = \bigcap_{j\in\mathbb{N}} \Lambda_{jp_0+n_0+m_0}.$$

Now using (9) we obtain

$$\Lambda = \bigcap_{j\in\mathbb{N}} f^{jp_0}(\Lambda_{n_0+m_0}) = \bigcap_{j\in\mathbb{N}} \bigcup_{A\in\mathcal{A}_{n_0+m_0}} f^{jp_0}(A). \tag{11}$$

Let $y \in \Lambda$. By equality (11), for all $j \in \mathbb{N}$ there exists an atom $A(j) \in \mathcal{A}_{n_0+m_0}$ such that $y \in f^{jp_0}(A(j))$. Therefore, since $\mathcal{A}_{n_0+m_0}$ is finite, it must exists $A \in \mathcal{A}_{n_0+m_0}$ and a subsequence $\{j_k\}_{k\in\mathbb{N}}$ of $\{jp_0\}_{j\in\mathbb{N}}$ such that $y \in f^{j_k}(A)$ for all $k \in \mathbb{N}$. Now, denote y^* the p_0-periodic point of f in A. Since y and y^* belong both to $f^{j_k}(A)$ for all $k \in \mathbb{N}$, we have

$$0 \leqslant d(y^*, y) \leqslant \operatorname{diam}(f^{j_k}(A)) \leqslant \lambda^{j_k}\operatorname{diam}(A) \qquad \forall\, k \in \mathbb{N},$$

and it follows that $y = y^*$. We conclude that any point of Λ is one of periodic points contained in each atom of $\mathcal{A}_{n_0+m_0}$. These points being unique in each of these atoms, Λ is finite. $\square$

Now we start our study of the remaining case $d(\Lambda, \Delta) = 0$. In the examples studied so far this lead the attractor to be a Cantor set. So, first of all we ask whether the total disconnection of Λ under the hypothesis of Theorem 2.10 extends to the general case $d(\Lambda, L) \geqslant 0$.

Theorem 2.12. *Let $f : X \mapsto X$ be a piecewise contracting map. If at least one of the four following hypothesis is satisfied:*

1) *X is a one–dimensional Riemannian manifold,*

2) *$f|_{X\setminus\Delta}$ is one–to–one,*

3) *f is atomically one–to–one,*

4) $\lim_{n\to\infty} (\#\mathcal{A}_n)\,\lambda^n = 0$,

then Λ is totally disconnected.

Proof. 1) Consider the contracting open pieces $X_1, X_2, \dots, X_N$. For each $i \in \{1, \dots, N\}$ we have

$$X_i = \bigcup_{j=1}^{+\infty} I_{i,j}$$

where $I_{i,j}$ is an open interval (a connected component of X_i). In the notation above we agree to denote $I_{i,j} = \emptyset$, if the open piece X_i has only a finite number $N_i \geqslant 1$ of connected components and $j > N_i$. Since the X_i are pairwise disjoint, and the different connected components of an open set are also pairwise disjoint, we deduce:

$$I_{i_1,j_1} \cap I_{i_2,j_2} = \emptyset \quad \text{if} \quad (i_1, j_1) \neq (i_2, j_2).$$

The space X being a compact one–dimensional manifold, and the open intervals $I_{i,j}$ being pairwise disjoint, we have for all $A \subset X$

$$\operatorname{diam}(A) = \operatorname{diam}((\bigcup_{i,j} A \cap I_{i,j}) \bigcup (A \setminus \bigcup_{i,j} I_{i,j})) \geqslant \operatorname{diam}(\bigcup_{i,j} A \cap I_{i,j}) \geqslant \sum_{i,j} \operatorname{diam}(A \cap I_{i,j}). \tag{12}$$

From the definition of atom, we obtain for all $n \geqslant 2$,

$$\sum_{A\in\mathcal{A}_n} \operatorname{diam}(A) = \sum_{A\in\mathcal{A}_{n-1}} \sum_{i=1}^{N} \operatorname{diam}(f(A \cap X_i)) \leqslant \sum_{A\in\mathcal{A}_{n-1}} \sum_{i=1}^{N} \lambda \operatorname{diam}(A \cap X_i) = \lambda \sum_{A\in\mathcal{A}_{n-1}} \sum_{i,j} \operatorname{diam}(A \cap I_{i,j}),$$

which with (12) leads to

$$\sum_{A\in\mathcal{A}_n} \operatorname{diam}(A) \leqslant \lambda \sum_{A\in\mathcal{A}_{n-1}} \operatorname{diam}(A) \leqslant \lambda^{n-1} \sum_{A\in\mathcal{A}_1} \operatorname{diam}(A) \qquad \forall\, n \geqslant 2.$$

Denote $l(\cdot)$ to the Lebesgue measure in X. Since the space X is a compact one-dimensional Riemannian manifold, we have $l(B) \leqslant \operatorname{diam}(B)$ for any Lebesgue measurable set $B \subset X$. Then, we deduce

$$l(\Lambda) \leqslant l(\Lambda_n) \leqslant \sum_{A \in \mathcal{A}_n} l(A) \leqslant \sum_{A \in \mathcal{A}_n} \operatorname{diam}(A) \leqslant \lambda^{n-1} \sum_{A \in \mathcal{A}_1} \operatorname{diam}(A) \qquad \forall\, n \geqslant 1,$$

and since $\lambda \in (0,1)$ and the number of atoms in $\mathcal{A}_1$ is finite, we have $l(\Lambda) = 0$. The attractor Λ has null Lebesgue measure and then, Λ can not contain open interval. In other words, it is totally disconnected.

2) If $f|_{X \setminus \Delta}$ is one–to–one, then the definition of atom implies the map f is atomically one–to–one.

To prove the thesis in the cases 2) and 3), we will prove the following assertion (A):

If $x \neq y \in \Lambda$, then there exists $n_0 \geqslant 1$ such that x and y belong to different connected components of Λ_{n_0}. (A)

Since $\Lambda \subset \Lambda_n$ for all $n \geqslant 1$, once we proved (A), it will follow that any two points $x \neq y \in \Lambda$ belongs to different connected components of Λ, as wanted.

3) Let us prove the assertion (A). As $x \neq y \in \Lambda$ there exists $n_0 \geqslant 1$ such that x and y belong to different atoms in $\mathcal{A}_{n_0}$. If not, x and y would belong to a same atom of generation arbitrarily large, and thus of diameter arbitrarily small, which contradicts $x \neq y$. As f is atomically one–to–one, the different atoms of $\mathcal{A}_{n_0}$ form a partition in closed sets of Λ_{n_0}. Therefore, a connected component of Λ_{n_0} cannot intersects two different atoms of $\mathcal{A}_{n_0}$. Since $x \neq y$ belongs to different atoms of $\mathcal{A}_{n_0}$, they belong to different connected components of Λ_{n_0}.

4) To prove the assertion (A), let us argue by contradiction. Suppose for all $n \geqslant 1$ the two points $x \neq y$ belong to the same connected component $C_n(x,y)$ of Λ_n, and denote N_n to the number of atoms of generation n that intersect $C_n(x,y)$. Then

$$d(x,y) \leqslant \operatorname{diam}(C_n(x,y)) \leqslant N_n \lambda^n \operatorname{diam}(X) \leqslant (\#\mathcal{A}_n)\lambda^n \operatorname{diam}(X) \qquad \forall n \geqslant 1.$$

Therefore $x = y$, which contradicts our assumption. □

Although we cannot exclude the possibility of an attractor containing a non–singleton connected components, in all the systems studied in the previous literature, at least one of the conditions of Theorem 2.12 holds and thus the attractor is totally disconnected. By the Cantor–Bendixson theorem [20], any closed subset of X can be written as the disjoint union of a perfect subset (a closed set without isolated point, called the *perfect kernel* of X) and a countable subset. Therefore, since Λ is compact, if it is uncountable and totally disconnected then its perfect kernel is non–empty and is a Cantor set.

In most examples $d(\Lambda, \Delta) \neq 0$, and the attractor is a finite set of periodic orbits. In the few examples where $d(\Lambda, \Delta) = 0$, the attractor has an uncountable transitive component which is a Cantor set (the perfect kernel of Λ) and possibly one or several additional transitive components which are finite (see [11] for instance). To sum up, the piecewise contracting systems that have been treated so far in the previous literature, in particular those introduced in [11] and defined by piecewise affine contraction with partition in orthotopes, have all shown the following specific characteristics:

A) The attractor is totally disconnected.

B) The attractor, the non wandering set and the limit set coincide: $\Lambda = \Omega = L$.

C) The attractor Λ is composed of a finite number of transitive components Λ^i, that is $\Lambda = \bigcup_{i=1}^n \Lambda^i$.

D) If a transitive component is infinite it is a Cantor set.

E) For each transitive component Λ^i of the attractor, there exists at least one continuous extension $f_i : \Lambda^i \to \Lambda^i$ of the map $f|_{\Lambda^i \setminus \Delta}$ such that the dynamical system (Λ^i, f_i) is transitive.

By Theorem 2.10, all these properties are trivially true if $d(\Lambda, \Delta) \neq 0$. The forthcoming Section 3 is a series of examples of piecewise contracting maps the attractor of which satisfies $d(\Lambda, \Delta) = 0$. For the sake of completeness of this section, we anticipate the general properties abstracted from these examples.

We could not find examples for which A) does not hold, and neither for which B) holds but C) does not. Nevertheless, our examples show that all the other properties are not proper to piecewise contracting maps.

First, we show that the condition $d(\Lambda, \Delta) = 0$ does not imply that the attractor contains a Cantor set. In Example 3.1, it is simply a finite set, as in the case $d(\Lambda, \Delta) \neq 0$, but not necessarily composed of periodic orbits, since E) does not hold. In the three examples of subsection 3.2, the attractor has an infinite transitive component, but it is countable and therefore D) fails. In Example 3.6 the attractor has a unique transitive component which is uncountable, but is not reduced to a Cantor set (the perfect kernel of Λ), since it also contains in addition a countable but infinite set (and D) fails again).

Second, the recurrence properties imposed by B) and C) do not generally hold. In Examples 3.3 and 3.4, we prove that $\Lambda \neq L$ and property B) fails. This implies that the attractor is not an union of transitive components, and C) fails also. In Example 3.3 we prove that the attractor may contain, wandering points in addition to infinitely many non recurrent points, in particular $\Lambda \neq \Omega$. On the other hand, in Example 3.4 the attractor Λ contains infinitely many non–recurrent points, but they are all non–wandering, and $\Omega \neq L$.

3 Examples and counter–examples

The purpose of this section is to study the questions raised at the end of Section 2 by means of two–dimensional examples where the continuity pieces have piecewise smooth borders (except in the example 3.6, for which one piece of the boarder is just a continuous arc). The examples are organized by increasing cardinality of the attractor.

3.1 Finite non–periodic attractor

We give an example of a piecewise contracting map for which the attractor Λ and the limit set L coincide, are finite, and are made of a single transitive component intersecting the discontinuity set Δ. In this example, there is no way to extend f on the discontinuity set in such a way that (Λ, f) becomes a transitive dynamical system.

Example 3.1. Fix $n \in \mathbb{N}$ and let $f_1 : [0,1]^2 \to [0,1]^2$ be the contracting affine map with fixed point at a point $(x_1, 0)$ defined by

$$f_1(x,y) = \lambda_1(x,y) + (1-\lambda_1)(x_1, 0),$$

where $x_1 \in [0, 1/2)$. The contraction rate $\lambda_1 \in (0,1)$ is chosen such that the n first pre–images by f_1 of the point $(1/2, 0)$ lie inside the segment $(1/2, 1)$[4]. For all $k \in \{1, \ldots, n\}$ denote $(a_k, 0)$ to the pre–image by f_1^k of the point $(1/2, 0)$. It is easy to check that

$$a_k = x_1 + \lambda_1^{-k}\left(\frac{1}{2} - x_1\right) \in (1/2, 1).$$

In brief, $a_0 := 1/2$ and $f_1(a_k, 0) = (a_{k-1}, 0) \in (1/2, 1) \times [0, 1]$ for all $k \in \{1, \ldots, n\}$.

Analogously, let us fix $m \in \mathbb{N}$ and let $f_2 : [0,1]^2 \to [0,1]^2$ be the contracting affine map defined by

$$f_2(x,y) = \lambda_2(x,y) + (1-\lambda_2)(x_2, 0),$$

where $x_2 \in (1/2, 1)$, and $\lambda_2 \in (0,1)$ is such that the m first pre–images of the point $(1/2, 0)$ lie in $(0, 1/2) \times [0, 1]$. For all $k \in \{1, \ldots, m\}$, let us denote $(b_k, 0)$ to the pre–image by f_2^k of the point $(1/2, 0)$. It is easy to check that

$$b_k = x_2 + \lambda_2^{-k}\left(x_2 - \frac{1}{2}\right) \in (0, 1/2).$$

---

[4] After a straightforward computation, one may choose

$$\lambda_1 \in \left(\left(\frac{1-x_1}{(1/2)-x_1}\right)^{-\frac{1}{n}}, \left(\frac{1-x_1}{(1/2)-x_1}\right)^{-\frac{1}{n+1}}\right] \subset (0,1).$$

In brief, $b_0 := 1/2$ and $f_2(b_k, 0) = (b_{k-1}, 0) \in (0, 1/2) \times [0, 1]$ for all $k \in \{1, \ldots, m\}$.

Let $f_3 : [0,1]^2 \to [0,1]^2$ be the following contracting affine map transforming the point $(1/2, 0)$ into $(a_n, 0)$, and the point $(0, 0)$ into $(1, 0)$:

$$f_3(x, y) = 2(1 - a_n)(1 - x, y) + (2a_n - 1, 0).$$

Then f_3 is a one–to–one contracting map from $[0, 1/2] \times [0, 1]$ into $[a_n, 1] \times [0, 1]$.

Finally let $f_4 : [0,1]^2 \to [0,1]^2$ be the contracting affine map transforming the point $(1/2, 0)$ into $(b_m, 0)$, and the point $(1, 0)$ into $(0, 0)$, defined by

$$f_4(x, y) = 2b_m(1 - x, y).$$

Then f_4 is a one–to–one contracting map from $[1/2, 1] \times [0, 1]$ into $[0, b_m] \times [0, 1]$.

Construct, as in Figure 1, the following open subsets of X:

$$X_1 = (a_1, 1] \times [0, 1], \quad X_2 = [0, b_1) \times [0, 1], \quad X_3 = (b_1, 1/2) \times [0, 1] \text{ and } X_4 = (1/2, a_1) \times [0, 1].$$

Notice that the point $(1/2, 0)$ belongs to $\partial X_3 \cap \partial X_4 \subset \Delta$.



Figure 1: Transitive attractor that is a ghost periodic orbit. The map f is an affine contraction of each one of the pieces of continuity $\{X_i\}_{i=1,\ldots,4}$.

Proposition 3.1. *Let $X = [0,1]^2$, and $f : X \to X$ such that $f|_{X_i} = f_i$ for all $i \in \{1, 2, 3, 4\}$ where the affine maps f_i and domains X_i are constructed as above. Then*

$$L = \{(x, 0) : x \in \{z_k\}_{k \in \mathbb{N}}\} = \Lambda,$$

where $\{z_k\}_{k \in \mathbb{N}}$ is the $(m + n + 2)$–periodic sequence given by:

$$\{z_k\}_{k=0,\ldots,m+n+1} = \{1/2, b_m, b_{m-1}, \ldots, b_1, 1/2, a_n, \ldots, a_1\} \quad \textit{and} \quad z_{k+m+n+2} = z_k \quad \textit{for all } k \in \mathbb{N},$$

and the attractor is transitive.

Remark 3.2. Although the sequence $\{(z_k,0)\}_{k\in\mathbb{N}}$ attracts any orbit of $(\tilde{X},f)$, it is neither an orbit of (X,f), nor an orbit of any of its possible continuous extensions. In particular, for most of the choices of f in Δ, the attractor is not invariant and despite being finite and transitive, it is not periodic. Nevertheless, if one would admit a multi–definition of f on Δ and would allow to choose, at each iteration, an image of a different continuous extension, a_n or b_m, of f on $(1/2,0)\in\Delta$, the attractor would be a periodic orbit for f. This example shows that the continuous extensions of f on Δ determines the attractor of $(\tilde{X},f)$.

Proof. Let $\lambda=\max\{\lambda_1,\lambda_2,2(1-a_n),2b_m\}$. Fix $(x,y)\in\tilde{X}$ with $x>1/2$. Compute the i_0 such that $0\leqslant i_0\leqslant n$ and $a_{i_0}<x<a_{i_0+1}$, defining $a_{n+1}=1$. Then

$$f^{i_0}(x,y)\in X_4,\quad f^{i_0+1}(x,y)=f_4(f^{i_0}(x,y))\in(0,b_m)\times[0,\lambda^{i_0+1}].$$

Therefore $f^{i_0+k}(x,y)$ belongs to an open rectangle D_k, constructed as the product of two open intervals of length λ^k, with the point $(z_k,0)$ at its boundary. Then $\lim_{k\to\infty}d(f^{i_0+k}(x,y),(z_k,0))=0$. Then for all $(x,y)\in\tilde{X}$ there exists $t_0\in\mathbb{N}$ such that

$$\lim_{k\to+\infty}d\left(f^{t_0+k}(x,y),(z_k,0)\right)=0.$$

A similar argument applies to $(x,y)\in\tilde{X}$ with $x<1/2$. It follows that

$$L=\{(x,0):x\in\{z_k\}_{k\in\mathbb{N}}\}.$$

After Definition 2.4, the arguments above also prove that $\Lambda_{m+n+2+k}$ is an union of less than $4(m+n+1+k)$ closed rectangles for all $k\in\mathbb{N}$, with sides not larger than λ^k, and with their respective boundaries intersecting the limit set L. Therefore: $\Lambda=\bigcap_{k\in\mathbb{N}}\Lambda_{m+n+2+k}=L$.

Finally, since $\omega(x,y)=L$ for all $(x,y)\in\tilde{X}$, Λ is transitive. □

3.2 Countably infinite attractor

In this subsection, we give three examples of piecewise contracting maps with countably infinite attractor.

The first and second of the following three examples, are defined in $X=[0,1]^2$. See Figure 2: left and right frames respectively. They are affine and contracting in the open sets

$$X_1=\{(x,y):y<x^2\}\quad\text{and}\quad X_2=\{(x,y):y>x^2\},\tag{13}$$

with the parabolic boundary $\Delta=\{(x,y):y=x^2\}$.

Example 3.2. Let $\lambda\in(0,1)$ and define on X the maps

$$f_1(x,y)=\lambda(x,y)\quad\text{and}\quad f_2(x,y)=(1-\lambda x,\lambda y).$$

See the left frame of Figure 2. Then we have the following:

Proposition 3.3. *Let $X=[0,1]^2$, and $f:X\to X$ such that $f|_{X_i}=f_i$ for $i=1,2$ where the affine maps f_i and the domains X_i are constructed as above. Then*

$$L=\overline{\{(\lambda^n,0):n\in\mathbb{N}\}}=\Lambda,$$

and the attractor is transitive.

Remark 3.4. The attractor is transitive according to Definition 2.5, but none of the infinitely many points of the attractor that belong to $\tilde{X}$ is recurrent.

Proof. Let $(x,y)\in\tilde{X}$ and let $(x_n,y_n)=f^n(x,y)$ for all $n\in\mathbb{N}$. Suppose $(x,y)\in X_1$. If $y=0$, then $f_1^n(x,y)=\lambda^n(x,0)\in X_1$ for any $n\in\mathbb{N}$. Therefore $f^n(x,y)=f_1^n(x,y)$ and $\omega(x,y)=\{(0,0)\}$. If $y\neq0$, then $y_n\neq0$ for all $n\in\mathbb{N}$ and there exists an increasing sequence $\{n_k\}_{k\in\mathbb{N}}$ such that $(x_{n_k},y_{n_k})\in X_2$ for all $k\in\mathbb{N}$. Moreover, by definition of X_2, we have $0<x_{n_k}<\sqrt{y_{n_k}}=\sqrt{\lambda^{n_k}y}$ which implies that



Figure 2: Countably infinite attractor. The example in the left frame is such that $L = \Omega = \Lambda$ with countable infinite limit set, while the one in the right frame is such that $L = \Omega \subsetneq \Lambda$ with finite limit set.

$\lim_{k\to\infty} f^{n_k}(x,y) = (0,0)$. We deduce that $\lim_{k\to\infty} f^{n_k+1}(x,y) = \lim_{k\to\infty} f_2(f^{n_k}(x,y)) = f_2(0,0) = (1,0)$ and it follows that

$$\lim_{k\to\infty} f^{n_k+1+n}(x,y) = f_1^n(1,0) = (\lambda^n, 0) \qquad \forall n \in \mathbb{N}.$$

In other words, $\omega(x,y) = \overline{\{(\lambda^n,0) : n \in \mathbb{N}\}}$. Now suppose $(x,y) \in X_2$. Then, $f^n(x,y) \in X_1$ for some $n \in \mathbb{N}$ and the proof is the same as before. Therefore, $L = \overline{\{(\lambda^n,0) : n \in \mathbb{N}\}}$ and is transitive.

After Theorem 2.9, to end this proof is enough to show that $\Lambda \subset L$. Note that for all $\epsilon > 0$ and for all $(x,y) \in \tilde{X}$ there exists $n \in \mathbb{N}$ such that $d(f^n(x,y),(0,0)) \leqslant \epsilon$. Therefore any point $(x,y) \in \Lambda$ is covered by a closed square, of side no larger than ϵ, intersecting L. Hence, $d((x,y),L) \leqslant \epsilon$ for all $\epsilon > 0$, and as L is compact we conclude that $(x,y) \in L$. $\square$

The second example is motivated by the Theorem 2.9. It shows that the limit set L and the attractor Λ may differ. The map differs from the previous example in its definition on X_2.

Example 3.3. Let $\lambda \in (0,1)$ and define on X the maps

$$f_1(x,y) = \lambda(x,y) \quad \text{and} \quad f_2(x,y) = \lambda(x,y) + (0, 1-\lambda).$$

See the right frame of Figure 2. Then we have the following:

Proposition 3.5. *Let $X = [0,1]^2$, and $f : X \to X$ such that $f|_{X_i} = f_i$ for $i = 1,2$ where the affine maps f_i and the domains X_i are constructed as above. Then*

$$L = \Omega = \{(0,0),(0,1)\} \qquad \textit{and} \qquad \Lambda = \overline{\{(0,1-\lambda^n) : n \in \mathbb{N}\}}.$$

Remark 3.6. The hypothesis $\Lambda \setminus L \subset \operatorname{int}(\tilde{X})$ of the last assertion in Theorem 2.9 holds in this example. Furthermore, as $\Lambda \setminus L \neq \emptyset$ and $L = \Omega$, the attractor Λ contains wandering points. Note that the attractor cannot be recurrent but has two transitive components (the points of L).

Proof. Let $(x,y) \in \tilde{X}$. Since the point $(0,1)$ is a global attractor for f_2 and $f(X_2) \subset X_2$, if $(x,y) \in X_2$ its limit set is $\omega(x,y) = \{(0,1)\}$. If $(x,y) \in X_1$ and $y = 0$ then $\omega(x,y) = \{(0,0)\}$, after the same argument in the proof of the proposition in the previous Example 3.2. If $y \neq 0$, then there exists $n \geqslant 1$ such that $f^n(x,y) \in X_2$ and therefore $\omega(x,y) = \{(0,1)\}$. It follows that $L = \{(0,0),(0,1)\}$.

Now we show that $\Lambda = \{(0, 1-\lambda^n) : n \in \mathbb{N}\} \cup \{(0,1)\}$. After Definition 2.4, $(x,y) \in \Lambda$ if and only if for all $n \geqslant 1$: $(x,y) \in A$ for some $A \in \mathcal{A}_n$. So $(x,y) \in \Lambda$ if and only if for all $n \geqslant 1$ there exists an admissible sequence $(i_1, \dots, i_n)$ of length n such that $(x,y) \in A_{i_1,\dots,i_n}$. Since $\overline{f(X_1)} \cap X_1 \neq \emptyset$, $\overline{f(X_1)} \cap X_2 \neq \emptyset$ and $\overline{f(X_2)} \subset X_2$, there are only three types of admissible sequences of length n: $(1,1,1,\dots,1)$, $(2,2,2,...,2)$ and $(1,1,...,1,2,2,...2)$. So, $\Lambda = \bigcup_{i=j}^{3} \Gamma_j$, where the subsets Γ_1, Γ_2 and Γ_3 are defined as follows:

– Γ_1 is formed by all the points $(x,y) \in X$ such that $x \in A_{1,1,1,\dots,1} \in \mathcal{A}_n$ for all $n \geqslant 1$.

– Γ_2 is formed by all the points $(x,y) \in X$ such that $x \in A_{2,2,2,\dots,2} \in \mathcal{A}_n$ for all $n \geqslant 1$.

– Γ_3 is formed by all the points $(x,y) \in X$ for which there exists $k \geqslant 1$ such that $(x,y) \in A_{2,2,2,\dots,2} \in \mathcal{A}_n$ for all $1 \leqslant n \leqslant k$, and $(x,y) \in A_{1,1,1,\dots,1,2,2,\dots,2} \in \mathcal{A}_n$ for all $n \geqslant k+1$, where exactly the last k symbols in the word $(1,1,1,\dots,1,2,2,\dots,2)$ of length n are "2". Note from the definition 2.4 of atom that Λ_3 is forward f_2–invariant.

Observe that $(0,0) \in \Gamma_1$. So, if $(x,y) \in \Lambda_1$ then $d((x,y),(0,0)) \leqslant \operatorname{diam}(A_{1,1,1,\dots,1}) \leqslant \max_{A \in \mathcal{A}_n} \operatorname{diam}(A) \leqslant \operatorname{diam}(X)\lambda^n$ for all $n \geqslant 1$. As $0 < \lambda < 1$, the last inequality implies that $(x,y) = (0,0)$. We deduce that $\Gamma_1 = \{(0,0)\}$. Analogously it is proved that $\Gamma_2 = \{(0,1)\}$. To prove that $\Gamma_3 \supset \{(0, 1-\lambda^m) : m \in \mathbb{N} \setminus \{0\}\}$, as Γ_3 is f_2–forward invariant, it is enough to prove that $(0, 1-\lambda) \in \Gamma_3$. The point $(0, 1-\lambda)$ belongs to A_2, to $A_{1,2}$, and to $A_{1,\dots,1,1,2} \in \mathcal{A}_n$ for all $n > 2$. After the definition of the subset Γ_3, we deduce that $(0, 1-\lambda) \in \Gamma_3$. To prove that $\Gamma_3 \subset \{(0,1-\lambda^m) : m \in \mathbb{N} \setminus \{0\}\}$, observe that, after the construction of the set Γ_3, for all $(x,y) \in \Gamma_3$ there exists $k \geqslant 1$ such that for all $n \geqslant k+1$, $(x,y) \in A_{1,1,1,\dots,1,2,\dots,2} = (F_2)^k \circ F_1^{n-k}(X) \subset f_2^k\left(\overline{X_2} \cap \overline{f_1^{n-k}(X_1)}\right)$, where F_i is the transformation defined in 2.4. Since $(0,0) \in \overline{X_2} \cap \overline{f_1^{n-k}(X_1)}$ for all $n \geqslant k+1$, we deduce that for all $\epsilon > 0$

$$d((x,y), f_2^k(B_\epsilon(0,0))) \leqslant \operatorname{diam}\left(f_2^k\left(\overline{X_2} \cap \overline{f_1^{n-k}(X_1)}\right)\right) < \lambda^k \operatorname{diam}\left(\overline{f_1^{n-k}(X_1)}\right) \leqslant \lambda^n \operatorname{diam}(X) \quad \forall\ n \geqslant k+1.$$

Taking $n \to +\infty$, we deduce that for all $\epsilon > 0$ $d((x,y), f_2^k(B_\epsilon(0,0))) = 0$, that is to say $(x,y) = \lim_{\epsilon \to 0} f_2^k(B_\epsilon(0,0)) = (0, 1-\lambda^k)$. Therefore, $\Lambda = \overline{\{(0,1-\lambda^n) : n \in \mathbb{N}\}}$.

Finally, from the above derivations we have that

$$\Lambda \setminus L = \{(0, 1-\lambda^n),\ n \in \mathbb{N} \setminus \{0\}\}.$$

The last set is contained in X_2, which is open in the relative topology of the space $X = [0,1]^2$. Besides, in this example $X_2 \subset \tilde{X}$. Then $\Lambda \setminus L \subset \operatorname{int}(\tilde{X})$. After Proposition 2.9 we have that $\Omega = L$. □

The third example is also motivated by Theorem 2.9 and shows an opposite case to the Example 3.3 where all the points of $\Lambda \setminus L$ are wandering. It shows that the limit set L and the non–wandering set Ω may differ.

Example 3.4. We construct a piecewise contracting map f in $X := [-1,1] \times [0,1]$ with three pieces of contraction:

$$X_1 := \{(x,y) \in X : 0 < x \leqslant 1, \quad 0 \leqslant y < x^2\},$$

$$X_2 := X_2^+ \ \cup\ \{(x,y) \in X : x = 0\}\ \cup\ X_2^-, \quad \text{where}$$

$$X_2^+ := \{(x,y) \in X : 0 < x \leqslant 1 \text{ and } \ y > x^2\} \ \text{ and } \ X_2^- := \{(x,y) \in X : -1 \leqslant x < 0 \text{ and } 0 \leqslant y < (x+1)^2\},$$

$$X_3 := \{(x,y) \in X : -1 \leqslant x \leqslant 0 \text{ and } (x+1)^2 < y \leqslant 1\}.$$

The compact set of discontinuities is $\Delta = \{(x,y) \in X : x \geqslant 0, y = x^2\} \bigcup \{(x,y) \in X : x \leqslant 0, y = (x+1)^2\}$. Let $0 < \lambda < 1$ and define on X the maps

$$f_1(x,y) = \lambda(x,y), \qquad f_2(x,y) = \lambda(x,y) + (0, 1-\lambda) \quad \text{and} \quad f_3(x,y) = \lambda(x,-y) + (1,\lambda).$$

See both frames of Figure 3. Then we have the following



Figure 3: Countably infinite attractor. In this example the limit set L is finite and $L \subsetneq \Omega = \Lambda$. In the left frame the half space has negative abscissa. The segments with zero abscissa in both frames are identified, and is plotted twice with separated frames for readability.

Proposition 3.7. *Let* $X = [-1, 1] \times [0, 1]$, *and* $f : X \to X$ *such that* $f|_{X_i} = f_i$ *for* $i = 1, 2, 3$ *where the affine maps* f_i *and the domains* X_i *are constructed as above. Then*

$$L = \{(0, 0), (0, 1)\} \qquad \text{and} \qquad \Omega = \Lambda = \overline{\{(0, 1 - \lambda^n) : n \in \mathbb{N}\}}.$$

Remark 3.8. Since $\Lambda \setminus L \not\subset \operatorname{int}(\tilde{X})$, the hypothesis of the last assertion of Theorem 2.9 does not hold, and the points of the non–empty set $\Lambda \setminus L = \Omega \setminus L$ are non–wandering. Besides, the thesis $L = \Omega$ of that theorem fails. Thus, it shows that the hypothesis of Theorem 2.9 is not redundant to imply that $L = \Omega$.

Proof. Let $(x, y) \in \tilde{X}$. For the same reason as in the previous Example 3.3, the ω-limit set of any point $(x, y) \in X_2^+ \bigcup \{(x, y) \in X : x = 0\}$ is $\omega(x, y) = \{(0, 1)\}$. The same proof of the proposition in the Example 3.3 also leads to $\omega(x, y) = \{(0, 0)\}$ if $(x, y) \in X_1$ and $y = 0$, and $\omega(x, y) = \{(0, 1)\}$ if $(x, y) \in X_1 \cap \tilde{X}$ and $y > 0$. On the other hand, if $(x, y) \in X_2^- \cap \tilde{X}$, there exists $n \geqslant 1$ such that $f^n(x, y) \in X_3$. Besides, if $(x, y) \in X_3 \cap \tilde{X}$, then $f(x, y) \in X_1 \cup X_2^+$. Therefore $\omega(x, y) \subset \{(0, 0), (0, 1)\}$ for all $x \in (X_2^- \cup X_3) \cap \tilde{X}$. Thus $L = \{(0, 0), (0, 1)\}$.

The proof of $\Lambda = \overline{\{(0, 1 - \lambda^n) : n \in \mathbb{N}\}}$ also follows the same arguments of the proof of the proposition in the previous Example 3.3.

After Theorem 2.9 we know that $L \subset \Omega \subset \Lambda$. As Ω is essentially forward invariant, to show that $\Omega = \overline{\{(0, 1 - \lambda^n) : n \in \mathbb{N}\}}$ it is enough to prove that the point $x_1 := (0, 1 - \lambda)$ belongs to Ω. For all sufficiently small $\epsilon > 0$, the ball $B_\epsilon(x_1)$ contains the segment $S = \{(-t, 1 - \lambda) : 0 < t < \epsilon\} \subset X_2^-$. Note that f_2 is an affine contraction with fixed point $(0, 1)$ that transforms horizontal lines to horizontal lines. Thus, the image of S by f_2 is also a horizontal segment with ordinate smaller than one. For all values of n large enough, denote $y_n = \{(-t_n, 1 - \lambda)\} \in S$ the point in S such that $f^i(y_n) \in X_2$ for all $i = 1, 2, \ldots, n-1$ and $f^n(y_n) \in \partial X_3 \cap \partial X_2 \subset \Delta$. After a straightforward computation it is not difficult to check that t_n is a strictly decreasing sequence of positive real numbers converging to zero. Consider for all n large enough the sub–segment $S_n = \{(-t, 1 - \lambda) : t_n < t < t_{n-1}\} \subset S$. By construction $f^n(S_n) = f_2^n(S_n) \subset X_3$ and so $d(f^n(S_n), (0, 1)) \leqslant \lambda^n$. Therefore

$$f^{n+1}(S_n) = f_3 \circ f_2^n(S_n) \subset X_1 \text{ and } d(f^{n+1}(S_n), (1, 0)) \leqslant \lambda^{n+1}. \tag{14}$$

For each n large enough, choose a natural number $m(n) \geqslant 1$ and a point $z_n \in f^{n+1}(S_n)$ such that $f^j(z_n) \in X_1$ for all $0 \leqslant j \leqslant m(n) - 1$ and $f^{m(n)}(z_n) \in X_2$. By construction $f^{m(n)}(z_n) = f_1^{m(n)}(z_n)$ and $f^{m(n)+1}(z_n) = f_2 \circ f_1^{m(n)}(z_n)$. The map f_1 is an affine contraction with fixed point $(0,0)$. So, after Equality (14), we deduce that $d(f^{m(n)}(z_n), (0,0)) \leqslant \lambda^{m(n)+n+1}$. Since f_2 is also an affine contraction, and $f_2(0,0) = (0, 1-\lambda) = x_1$ we deduce that

$$d(f^{m(n)+1}(z_n), x_1) = d(f_2 \circ f_1^{m(n)}(z_n), x_1) \leqslant \lambda^{m(n)+n+2}.$$

We conclude that for any n large enough, the point $w_n = f^{m(n)+1}(z_n)$ belongs to the ball $B_\epsilon(x_1)$. By construction, $z_n \in f^{n+1}(S_n)$ and $S_n \subset B_\epsilon(x_1)$, so $w_n \in f^{m(n)+n+2}(B_\epsilon(x_1))$. Finally, denoting $k_n = m(n) + n + 2$ we deduce that

$$w_n \in B_\epsilon(x_1) \bigcap f^{k_n}(B_\epsilon(x_1)) \neq \emptyset.$$

As the proof above holds for any arbitrarily small $\epsilon > 0$, after Definition 2.3, the point x_1 belongs to the non–wandering set Ω.

Finally, if the last assertion were false then, due to the part b) of Theorem 2.9, the non–wandering set Ω would be equal to the limit set L. But, as already proved, $L \neq \Omega$. Therefore $\Lambda(f) \setminus L(f) \not\subset \operatorname{int}(\tilde{X})$. ☐

3.3 Uncountable limit set

There are known piecewise contracting maps for which the attractor Λ is uncountably infinite, totally disconnected and has several transitive components, each being either a finite or a perfect set. The simplest among those examples can be found in [11] and consists of piecewise affine maps on the plane with a partition in rectangles. In this case the attractor is the union of one transitive component which is a Cantor set (the closure of a quasi–periodic orbit) and zero, one or several other transitive components, which are finite sets (corresponding to as many periodic orbits) [11, Theorems 4.1 and 4.2].

Here we give the Example 3.6 of a piecewise contracting map in the plane exhibiting an attractor Λ which is uncountably infinite and totally disconnected, which has a unique transitive component but has two invariant subsets: a perfect kernel K which is a Cantor set, and $C = \Lambda \setminus K$ which is countably infinite.

For the construction of this two dimensional piecewise contracting example, we need to recall the existence of another example consisting of a piecewise contracting map h on the interval $[0,1] \subset \mathbb{R}$, taken from [11].

Example 3.5. Let $X = [0,1]$, $X_1 := (\tau, 1]$ and $X_2 := [0, \tau)$ with $\tau \in (0,1)$. Let $\alpha \in (0,1)$ and define on X the maps:

$$h_1 = \alpha x \quad \text{and} \quad h_2 = \alpha x + (1-\alpha).$$

Then we have the following:

Proposition 3.9. *Let $X = [0,1]$ and $h : X \to X$ such that $h|_{X_i} = h_i$ for $i = 1, 2$ where the affine maps h_i and the domains X_i are constructed as above. Then for all $\tau \in (0,1) \setminus \{1/2\}$ and $\nu \in (0,1) \cap (\mathbb{R} \setminus \mathbb{Q})$ there exists $\alpha \in (0,1)$ such that*

$$T(\alpha, \nu) = \tau,$$

where $T : (0,1) \times \big\{\{\mathbb{R} \setminus \mathbb{Q}\} \cap [0,1]\big\} \to (0,1)$ is defined by

$$T(\alpha, \nu) := 1 - \frac{(1-\alpha)^2}{\alpha} \sum_{k=0}^{\infty} \alpha^k \lfloor \nu(k+1) \rfloor, \tag{15}$$

and for all $(\alpha, \nu) \in (0,1) \times \big\{\{\mathbb{R} \setminus \mathbb{Q}\} \cap [0,1]\big\}$ the set

$$K = \overline{\big\{h^n(T(\alpha,\nu))\big\}_{n \in \mathbb{N}}}$$

is a Cantor set that satisfies: $0 \leqslant \min K < T(\alpha, \nu) < \max K \leqslant 1$. Furthermore,

$$L \ = \ K \ = \ \Lambda,$$

and the attractor is transitive.

In the Formula (15), $\lfloor x \rfloor$ denotes the largest integer that is not larger than x.

Proof. A corrolary of [11, Theorem 6.1]. $\square$



Figure 4: The attractor is the union of a Cantor set and a countably infinite set. In this example $L=\Omega=\Lambda$.

In the following Example 3.6, using the above property of $([0,1],h)$, we construct a piecewise contracting map $f: X \mapsto X$, where $X=[0,2]\times[0,1]\subset\mathbb{R}^2$ has five pieces of contraction $\{X_i\}_{i=1,\dots,5}$, as in Figure 4. The construction follows five steps. First, applying Theorem 3.9, we choose adequate values of the two real parameters $\tau\in(0,1)$ and $\alpha\in(0,1)$ for the one–dimensional map h according to Formula (15), and we distinguish from the properties of its attractor two points to be used in the following steps of the construction of (X,f). Second, we construct an adequate continuous arc to become the shared boundary $\partial X_1 \bigcap \partial X_2 = \Delta_{12}$ of the two open pieces X_1 and X_2, as in Figure 4. Third, we define five open pieces X_1, X_2, X_3, X_4 and X_5 to be the pieces of contraction of the piecewise contracting map $f: X\mapsto X$. Fourth, we construct a two–dimensional map g, such that $g|_{[0,1]}=h$, with h as above, and we study its properties in relation to the boundary Δ_{12} that was constructed in the second step. Fifth and last, we construct five contracting homeomorphisms $\{f_i\}_{i=1,\dots,5}$ from $\mathbb{R}^2$ to $\mathbb{R}^2$, such that all of them, except f_4 is an affinity, and each f_i transforms the respective piece X_i into an image inside X. So we define a piecewise contracting map $f: X\mapsto X$ such that $f|_{X_i}=f_i$ for all $i=1,\dots,5$. The Proposition 3.12 states the asymptotic dynamical properties of the map f this way constructed. We show that its attractor is transitive and formed by two mutually disjoint compact subsets: a Cantor set and a countably infinite set.

Example 3.6. First step. Fix arbitrary values for $\alpha\in(0,1)$ and $\nu\in(0,1)\cap\{\mathbb{R}\setminus\mathbb{Q}\}$ and find $\tau\in(0,1)$ satisfying the thesis of Theorem 3.9 using (15). Using such α and τ, construct the piecewise affine map $h:[0,1]\mapsto[0,1]$ defined in Example (3.5), and denote

$$x_s := \min K,$$

where K is the Cantor set characterized in Theorem 3.9. In Figure 4 we have drawn some points of the set K, contained in the segment $\{(x,y)\in\mathbb{R}^2: 0\leqslant x\leqslant 1,\ y=0\}$. We now choose a point $x_e\in K$ such that:

$$\tau < x_e < \max K \quad \text{and} \quad \forall\, n\in\mathbb{N}:\ x_s \neq h^n(x_e) \neq \tau.$$

The existence of such a point $x_e\in K$ is provided by Theorem 3.9. Indeed, the attractor K intersects the two open sub–intervals, and as K is perfect and compact there exist uncountably many points of K that are larger than τ. Among those points there exists uncountably many that are not in the countable set formed with all the pre–images by $h, h^2, \dots, h^n, \dots$ of the points x_s and τ.

Second step. Consider the strictly increasing sequence $\{n_i\}_{i\geqslant 1}$ of natural numbers defined by:

$$n_1 := \min\{n \geqslant 1 : h^n(x_e) < \tau\}, \quad \text{and} \quad n_{i+1} := \min\{n > n_i : h^n(x_e) < h^{n_i}(x_e)\} \ \forall\, i > 1,$$

Such sequence $\{n_i\}_{1\geqslant 1}$ exists because $x_e \in K$ was chosen such that its forward orbit by h does not include the point $x_s = \min K$. So for any $n \geqslant 0$, there exists points of K that are at the left of $f^n(x_e)$ in the interval $[0,1]$. After Theorem 3.9, $K = \omega_h(x_e)$. So, the forward orbit of x_e accumulates to any point of K, in particular to those points of K that are at the left of τ and at the left of $f^n(x_e)$. Notice that by construction of the sequence $\{n_i\}_{i\geqslant 1}$, the sequence of points $\{h^{n_i}(x_e)\}_{i\in\mathbb{N}}$ is strictly decreasing in the interval $[0,1]$.

Choose a real number ρ such that $0 < \rho < \alpha < 1$. Construct a piecewise affine and continuous function $\phi : [x_s, \tau] \to [0,1]$ such that it is increasing and besides:

$$\phi(x_s) = 0, \quad \phi(\tau) = 1 \ \text{ and } \ \phi(h^{n_i}(x_e)) = \rho^{n_i} \quad \forall\, i \in \mathbb{N}. \tag{16}$$

Denote the graph of ϕ:

$$\Delta_{12} = \{(x, \phi(x)) : x_s \leqslant x \leqslant \tau\},$$

and define $\partial X_1 \cap \partial X_2 = \Delta_{12}$

Lemma 3.10. *The real function ϕ, whose graph Δ_{12} is constructed as above, satisfies the following property: For any fixed positive value of $y_0 \in (0,1)$, there exists a natural number $n_0 \geqslant 1$ such that*

$$h^{n_0}(x_e) \in (x_s, \tau) \ \text{ and } \ \phi(h^{n_0}(x_e)) < \alpha^{n_0} y_0.$$

Besides, any sequence of positive real numbers $\{y_j\}_{j>0}$ such that $\lim_{j\to+\infty}(\log y_j)/j \leqslant \log\alpha$ satisfies:

$$\lim_{j\to+\infty} \min\{n \geqslant 1 : \ h^n(x_e) \in (x_s, \tau) \ \text{ and } \ \phi(h^n(x_e)) < \alpha^n\, y_j\} = +\infty.$$

Proof. The first assertion is proved by contradiction. If for all $n \geqslant 1$ such that $x_s < h^n(x_e) < \tau$ we had $\alpha^n\, y_0 \leqslant \phi(h^n(x_e)\,)$, then by definition of ϕ in Equation (16) we would obtain: $\alpha^{n_i}\, y_0 \leqslant \phi(h^{n_i}(x_e)\,) = \rho^{n_i}$ $\forall\, i \in \mathbb{N}$. Therefore $y_0 \leqslant (\rho/\alpha)^{n_i}$. Since $0 < \rho < \alpha$, taking $i \to +\infty$ we get $y_0 \leqslant 0$, contradicting our hypothesis $0 < y_0$.

The second assertion is proved as follows. Fix $j \in \mathbb{N}\setminus\{0\}$ and take $y_j > 0$. Denote $n(y_j) \geqslant 1$ the minimum $n \in \mathbb{N}$ satisfying the three inequalities in the last assertion of this lemma. Then

$$\phi(h^{n(y_j)}(x_e)\,) = \rho^{n(y_j)} < \alpha^{n(y_j)} y_j \quad \Rightarrow \quad \frac{1}{y_j} < \left(\frac{\alpha}{\rho}\right)^{n(y_j)}.$$

Thus

$$\frac{-\log y_j}{j} < \frac{n(y_j)}{j}\,(\log\alpha - \log\rho) \quad \forall j > 0.$$

Assume that $\lim_{j\to+\infty}(-\log y_j)/j \geqslant -\log\alpha > 0$. Then, from the definition of limit, for all $\epsilon > 0$ there exists $j_0 \in \mathbb{N}\setminus\{0\}$ such that $-(\log y_j)/j > -\log\alpha - \epsilon$ for all $j \geqslant j_0$. Therefore

$$\frac{-\log\alpha - \epsilon}{\log\alpha - \log\rho} < \frac{-(\log y_j)/j}{\log\alpha - \log\rho} < \frac{n(y_j)}{j} \quad \forall\, j \geqslant j_0.$$

Choosing $\epsilon = -(\log\alpha)/2 > 0$ and denoting $C = (-(\log\alpha)/2)\,/\,(\log\alpha - \log\rho)$, (note that $C > 0$, because $0 < \rho < \alpha < 1$), we conclude that $n(y_j) > jC$ for all $j \geqslant 0$, and thus $\lim_{j\to+\infty} n(y_j) = +\infty$. □

Third Step. Construct the topological partition of the compact rectangle $X := [0,2]\times[0,1] = \bigcup_{i=1}^{5} \overline{X}_i$, where $\{X_i\}_{1\leqslant i\leqslant 5}$ is the family of the following five open sets, as shown in Figure 4:

$$\begin{aligned}
X_1 &= \{(x,y) : x_s \leqslant x < \tau, \ \ \phi(x) < y \leqslant 1\} \ \cup \ \{(x,y) : 0 \leqslant x < x_s, \ \ 0 \leqslant y \leqslant 1\},\\
X_2 &= \{(x,y) : x_s < x < \tau, \ \ 0 \leqslant y < \phi(x)\},\\
X_3 &= \{(x,y) : \ \tau < x < 1, \ \ 0 \leqslant y \leqslant 1\},\\
X_4 &= \{(x,y) : \ 1 < x < 2, \ \ (x-1)^2 < y \leqslant 1\},\\
X_5 &= \{(x,y) : \ 1 < x \leqslant 2, \ \ 0 \leqslant y < (x-1)^2\}.
\end{aligned}$$

Notice that $X_1 \cup X_2 \cup X_3 = [0,1]^2$, and that by definition of X_2 and X_3 the point $(\tau, 0)$ belongs to $\Delta = \bigcup_{i \neq j} \partial X_i \cap \partial X_j$.

Fourth Step. Define the map $g : [0,1]^2 \to [0,1]^2$ as follows:

$$g(x,y) := (h(x),\ \alpha y) \text{ such that } x \neq \tau \text{ with } \alpha \text{ and } \tau \text{ as in Example 3.5.} \tag{17}$$

After Theorem 3.9, it follows that the attractor of g and its limit set are the Cantor set

$$K \times \{0\}.$$

Furthermore, after Lemma 3.10, for any fixed $y_0 \in (0,1)$ there exists a natural number $n_0 \geqslant 1$ such that

$$g^{n_0}(x_e, y_0) = (h^{n_0}(x_e), \alpha^{n_0} y_0) \in \{(x,y) : x_s < x < \tau,\ \phi(x) < y\} =: \check{X}_1 \subset X_1. \tag{18}$$

The following lemma will be fundamental to construct, in the next and last step, the piecewise contracting map $f : X \mapsto X$ with the desired properties.

Lemma 3.11. *There exists an open set $\mathcal{D} \subset X_3$, such that $(x_e, 0) \in \partial\mathcal{D}$, satisfying the following properties:*

a) *For any point $(x,y) \in \mathcal{D}$ there exists a natural number $n_0 \geqslant 1$, such that*

$$g^{n_0}(x,y) \in X_1,$$

where g is defined in Equality (17).

b) *For all $(x,y) \in \mathcal{D}$ and for all j such that $0 \leqslant j \leqslant \min\{n \in \mathbb{N} : \ g^n(x,y) \in X_1\}$:*

$$|g^j(x,y) - g^j(x_e, y)| \leqslant \alpha^j$$

c) *For any sequence of points $(x_j, y_j) \in \mathcal{D}$:*

$$\textit{If } \lim_{j \to +\infty} \frac{\log y_j}{j} \leqslant \log \alpha, \quad \textit{then } \lim_{j \to +\infty} m_j = +\infty, \ \textit{where } m_j := \min\{n \in \mathbb{N} : \ g^n(x_j, y_j) \in X_1\}.$$

d) *With the hypothesis of part* c), $\lim_{j \to +\infty} g^{m_j}(x_j, y_j) = (x_s, 0)$.

Proof. a): Choose any positive value of $y_0 \in (0,1)$. Then $(x_e, y_0) \in X_3 \cap \{(x,y) \in X : y < 1\}$. Recall that the point $x_e \in K \subset [0,1]$ was chosen in such a way that $h^n(x_e) \neq \tau$ for all $n \geqslant 0$. Since $g(x,y)$ is continuous if $x \neq \tau$, we deduce that the map g^n, for any $n \geqslant 1$, is defined and continuous at the point (x_e, y_0). Besides, after the assertion in Formula (18), there exists a minimum $n_0 \geqslant 1$ such that $g^{n_0}(x_e, y_0) \in \check{X}_1 \subset X_1$, and the sets $\check{X}_1$ and $X_3 \cap \{(x,y) \in X : y < 1\}$ are open in $\mathbb{R}^2$. Therefore, there exists an open ball

$$B(x_e, y_0) \subset X_3 \cap \{(x,y) \in X : y < 1\},$$

with center at the point (x_e, y_0) and radius, say, $\epsilon_0 > 0$ (depending on y_0), such that

$$g^{n_0}(B(x_e, y_0)) \subset \check{X}_1 \subset X_1.$$

Since g^n is continuous at (x_e, y_0) for all $n \geqslant 0$, and since $g^i(x_e, y_0) \in X_2 \cup X_3$ for all $i = 0, \dots, n_0 - 1$, the radius $\epsilon_0 > 0$ of the ball $B(y_0)$ can be chosen small enough, such that

$$\forall i = 0, \dots, n_0 - 1, \quad g^i(B(y_0)) \in X_2 \cup X_3. \tag{19}$$

Define the set

$$\mathcal{D} = \bigcup_{0 < y_0 < 1} B(y_0). \tag{20}$$

This set $\mathcal{D}$ is open because it is the union of open balls. The inclusion $\mathcal{D} \subset X_3$ is immediate recalling that all the balls $B(y_0)$ were constructed being contained in X_3. Besides, $(x_e, 0) \in \partial\mathcal{D}$ because $(x_e, y_0) \in B(y_0) \subset \mathcal{D}$ for all $0 < y_0 < 1$.

b): After Assertion (19), for all $(x, y) \in \mathcal{D}$, either $g^j(x, y) \in g^j(B(y_0)) \subset X_2$ or it is contained in X_3 for all $j = 1, 2, \ldots, n_0 - 1$ (because X_2 and X_3 are open sets, and thus $g^j(B(y_0))$ is the continuous image of a ball, and thus it is connected.) Since g^j is contracting with rate α in each open piece X_2 and X_3, the points $g^j(x, y) = (h^j(x), \alpha^j\, y)$ and $g^j(x_e, y_0) = h^j(x_e), \alpha^j\, y_0)$ are at distance smaller that α^j. Then $|h^j(x) - h^j(x_e)| \leqslant \alpha^j$. But the distance between $g^j(x, y)$ and $g^j(x_e, y) = (h^j(x_e), \alpha^j\, y)$ is exactly $|h^j(x) - h^j(x_e)|$.

c): Take any sequence $\{(x_j, y_j)\}_{j\geqslant 1}$ in $\mathcal{D}$ such that $\lim(\log y_j)/j \leqslant \log\alpha$. After Equality (20) and since $(x_j, y_j) \in \mathcal{D}$, there exists a ball $B_j = B(x_e, y_0(j))$ centered at the point $(x_e, y_0(j))$, such that $(x_j, y_j) \in B_j$ and such that for some natural number $n_0(j) \geqslant 1$ the center satisfies $g^{n_0(j)}(x_e, y_0(j)) \in X_1$. Then one has $g^{n_0(j)}(B_j) \subset X_1$, and in particular $g^{n_0(j)}(x_j, y_j) \in X_1$. Since m_j is the minimum natural number such that $g^{m_j}(x_j, y_j) \in X_1$ we deduce that:

$$m_j \leqslant n_0(j).$$

Now we prove that $m_j \geqslant n_0(j)$. By contradiction, if $m_j < n_0(j)$, we can apply Inequality (19): $f^{m_j}(B_j) \in X_2 \cup X_3$. So, in particular $g^{m_j}(x_j, y_j) \in X_2 \cup X_3$ and so $g^{m_j}(x_j, y_j) \notin X_1$, contradicting the construction of the natural number m_j. Then

$$m_j = n_0(j) \ \ \forall\, j \geqslant 1.$$

By hypothesis $\lim(-\log y_j)/j \leqslant \log\alpha$, so we can apply Lemma 3.10 to obtain

$$\lim_{j\to+\infty} m_j = \lim_{j\to+\infty} n_0(j) = +\infty.$$

d): From the definition of x_s, x_e, g given by Formula (17) implying that $g^n(x_e, 0) \in K \times \{0\}$ for all $n \geqslant 0$, and using the item b) of the Lemma, the following inequalities hold:

$$\operatorname{dist}(\ g^{m_j}(x_j, y_j),\ K \times \{0\}\ ) \leqslant \operatorname{dist}(\ g^{m_j}(x_j, y_j),\ g^{m_j}(x_e, 0)\ ) \leqslant$$

$$\leqslant \operatorname{dist}(\ g^{m_j}(x_j, y_j),\ g^{m_j}(x_e, y_j)\ )\ +\ \operatorname{dist}(\ g^{m_j}(x_e, y_j),\ g^{m_j}(x_e, 0)\ ) \leqslant \alpha^j + \alpha^j\, y_j \leqslant 2\alpha^j \ \ \forall\, j \geqslant 1.$$

Therefore, since $\alpha \in (0, 1)$ we obtain that $\lim_{j\to+\infty} \operatorname{dist}(\ g^{m_j}((x_j, y_j),\ K \times 0\ ) = 0$. Therefore any limit point of the sequence $\{g^{m_j}(x_j, y_j)\}_{j\geqslant 1}$ belongs to $K \times 0$. Besides, from the definition of m_j, the sequence $\{g^{m_j}(x_j, y_j)\}_{j\geqslant 1}$ is contained in X_1, and all its limit points are contained in $\overline{X_1}$. So, they are contained in $\overline{X_1} \cap (K \times \{0\}) = \{(x_s, 0)\}$. We conclude that the sequence $\{g^{m_j}(x_j, y_j)\}_{j\geqslant 1}$ is convergent, and that its limit is the point $(x_s, 0)$. □

Fifth Step. Construct five contracting invertible maps on the plane.

– $f_1 : \mathbb{R}^2 \to \mathbb{R}^2$ such that $f_1(x, y) = \lambda_1(x, y) + (\beta_1, 0)$, with any choice of the parameter β_1 such that $1 < \beta_1 < 2$, and with the contracting rate λ_1 chosen such that $0 < \lambda_1 < \min\left\{(\beta_1 - 1)^2, 2 - \beta_1/\tau\right\}$, where $0 < \tau < 1$ is the parameter fixed in the First Step. Then f_1 maps X_1 into X_5 (See Figure 4).

– $f_2 : \mathbb{R}^2 \to \mathbb{R}^2$ defined by $f_2(x, y) = \alpha(x, y) + (1 - \alpha, 0)$, where $0 < \alpha < 1$ is the real number of Proposition 3.9. Recalling formula (15), the two–dimensional affinity f_2 satisfies

$$f_2(x, y) = (h(x), \alpha y) = g(x, y) \ \text{ if } 0 \leqslant x < \tau,$$

where g is defined in Formula (17). Note that $(1, 0)$ is the unique fixed point of f_2 which is a contracting homothecy, and thus f_2 maps X_2 into $\overline{X_2} \cup \overline{X_3}$ in such a way that the iterates of the points of X_2 remaining in X_2 approach the Cantor set $K \times \{0\}$.

– $f_3 : \mathbb{R}^2 \to \mathbb{R}^2$ defined by $f_3(x, y) = \alpha(x, y)$. Note that

$$f_3(x, y) = (h(x), \alpha y) = g(x, y) \ \text{ if } \tau < x < 1.$$

Besides, $(0, 0)$ is the unique fixed point of f_3 which is a contracting homothecy, and thus it maps X_3 into $\overline{X_1} \cup \overline{X_2} \cup \overline{X_3}$ in such a way that the iterates of the points of X_3 approach the Cantor set $K \times \{0\}$ while they remain in X_3.

– $f_4 : \mathbb{R}^2 \to \mathbb{R}^2$, being any contracting homeomorphism, such that $f_4(1,0) = (x_e, 0)$, $f_4(X_4) \subset \mathcal{D}$, where $\mathcal{D}$ is the open set of Lemma 3.11, and if $(\bar{x}, \bar{y}) = f_4(x,y)$ then $\bar{y}$ is independent of x for all $(x,y) \in \mathbb{R}^2$, and for all $x \in [0,1]$, $f_4(x,y) > f_4(x,y')$ if $y > y'$ for all $y, y' \in [0,1]$. The map f_4 has the following property that will be used in the proof of Proposition 3.12 to show non–trivial asymptotic dynamics of the constructed piecewise contracting map f: If a sequence of points $\mathcal{O} = \{(x_n, y_n)\}_{n \geqslant 0}$ (which will be later an orbit of a dynamical system to be defined on X) visits infinitely many times the open piece X_4, in such a way that for all $i \geqslant 2$ the i–th visit $(x_{n_i}, y_{n_i}) \in X_4$ satisfies $y_{n_i} < y_{n_{i-1}}$ and $f_4(x_{n_i}, y_{n_i}) = (x_{n_i+1}, y_{n_i+1})$, then the sequence $\mathcal{O}$ also visits infinitely many times the open piece X_3 in the points $(x_{n_i+1}, y_{n_i+1}) \in \mathcal{D} \subset X_3$ in such a way that $y_{n_i+1} < y_{n_{i-1}+1}$ for all $i \geqslant 1$.

– $f_5 : X \to X$ defined by $f_5(x,y) = \lambda_5(x,y) + (1-\lambda_5, 0)$ with any choice of the contracting rate $\lambda_5 \in (0,1)$.

Finally, take the following piecewise contracting map:

$$f : X = [0,2] \times [0,1] \mapsto X \text{ such that } f|_{X_i} = f_i|_{X_i} \text{ for } i = 1,2,3,4,5.$$

It is immediate, after the construction above, that f satisfies the conditions 1) and 2) of Definition 2.1 of a piecewise contracting map. Besides, by construction, for every $1 \leqslant i \leqslant 5$ the transformation f_i is a homeomorphism on the plane. Then, after Theorem 2.8, the set $\tilde{X} = \bigcap_{n \geqslant 0} f^{-n}(X \setminus \Delta)$ is dense in X and the condition 3) of Definition 2.1 holds.

Now, we state and prove the following result about the asymptotic properties of f:

Proposition 3.12. *Let $X = [0,2] \times [0,1]$, and let $f : X \to X$ such that $f|_{X_i} = f_i$ for all $i \in \{1,2,3,4,5\}$ where the affine maps f_i and the domains X_i are defined above. Then:*

$$L = \mathcal{K} \cup \mathcal{S} = \Omega = \Lambda,$$

where the totally disconnected perfect kernel $\mathcal{K}$ is a Cantor set, $\mathcal{S}$ is a countably infinite set, and the attractor has is transitive.

To prove the proposition we need two lemmas about the set $\mathcal{K} \cup \mathcal{S}$ characterized below. Denote

$$\mathcal{K} := K \times \{0\},$$

where $K \subset [0,1]$ is the Cantor set attractor of the map h in Proposition 3.9. Recall that h is the map defined in Example (3.5), extended to a bi–dimensional map g in Formula (17). This map g, as observed in the fourth step, has $\mathcal{K} \subset [0,1] \times \{0\}$ as attractor and as limit set. Besides, in the fifth step of the construction of f such map g, when restricted to X_2 and X_3, coincides with f_2 and f_3, respectively. Thus g coincides with f in $X_2 \cup X_3$. So, we deduce that if $(x,y) \in X$ is such that for all arbitrary large values of n the iterates $f^n(x,y) \in X_2 \cup X_3$ then $\omega(x,y) = \mathcal{K}$. In particular,

$$\omega(x_e, 0) = \mathcal{K}, \tag{21}$$

where $(x_e, 0) \in X_3$ is the point chosen in the first step of the construction of f (see Figure 4). Besides $(x_e, 0) \in \tilde{X}$ because $f^n(x_e, 0) = g^n(x_e, 0) = (h^n(x_e), 0) \in X_2 \cup X_3$ for all $n \geqslant 0$. (Recall the construction of the pieces X_2 and X_3 in the fourth step above, and the choice of $x_e \in K$, in the first step, such that $h^n(x_e) \neq x_s, \tau$ for all $n \geqslant 0$.) So

$$\mathcal{K} \subset \overline{\bigcup_{(x,y) \in \tilde{X}} \omega(x,y)} = L.$$

Denote

$$\mathcal{S} := \{(1,0)\} \ \cup \ \big\{(1 + \lambda_5^k(\lambda_1 x_s + \beta_1 - 1), \ \ 0)\big\}_{k \geqslant 0} \ \subset [1,2] \times \{0\},$$

where $x_s = \min K$ (as defined in the first step), and $0 < \lambda_1, \lambda_5 < 1$ and $1 < \beta_1 < 2$ are real parameters used in the fifth step of the construction of f, when defining the affinities f_1 and f_5. It is standard to check, using the respective formulae of the affinities $f_1, f_5 : \mathbb{R}^2 \mapsto \mathbb{R}^2$, that:

$$\mathcal{S} = \{(1,0)\} \cup \{f_5^k \circ f_1(x_s, 0)\}_{k \geqslant 0} = \overline{\{f_5^k \circ f_1(x_s, 0)\}_{k \geqslant 0}}. \tag{22}$$

The set $\mathcal{K} \cup \mathcal{S}$ is, by construction, the union of the totally disconnected perfect kernel $\mathcal{K}$, which is a Cantor set in $X = [0,2] \times [0,1]$, plus a countably infinite set $\mathcal{S}$. We shall prove that this union is the attractor Λ of the piecewise contracting map f.

Lemma 3.13.

$$\Lambda \subset \mathcal{K} \ \cup \ \mathcal{S} \tag{23}$$

Proof. From Definition 2.4, a point (x,y) belongs to the attractor Λ if and only if for all $n \geqslant 1$ there exists an atom $A \in \mathcal{A}_n$ such that $(x,y) \in A$. So, to study the attractor, we study the admissibility of itineraries for points in the attractor.

Let $0 < \lambda < 1$ be the maximum contracting rate of f in its five pieces of contraction, and observe that the iterate $(\bar{x}, \bar{y}) = f(x,y)$ of any point $(x,y) \in X \setminus \Delta$ satisfies $0 \leqslant \bar{y} \leqslant \lambda y$. Therefore, any atom A of generation $n \geqslant 1$ satisfies $A \in \{(x,y) \in X : 0 \leqslant y \leqslant \lambda^n\}$. Then,

$$\bigcup_{A \in \mathcal{A}_n} A = \Lambda_n \subset \{(x,y) \in X : 0 \leqslant y \leqslant \lambda^n\} \ \ \forall \ n \geqslant 1,$$

and thus

$$\Lambda = \bigcup_{n \geqslant 1} \Lambda_n \ \subset \ \bigcup_{n \geqslant 1} \{(x,y) \in X : 0 \leqslant y \leqslant \lambda^n\} = \{(x,y) \in X : y = 0\}.$$

Let $n \geqslant 1$ and choose any atom $A \in \mathcal{A}_n$. Denote its admissible n–itinerary as $I_n(A) = (i_1, i_2, \ldots, i_n)$. Considering all the admissible n-itineraries, we obtain the four following cases:

(i) If $i_n = 1$, then $i_{n-1} \in \{2,3\}$, because the only pieces X_i such that $\overline{f(X_i)} \cap X_1 \neq \emptyset$ are X_2 or X_3. So, if a point $(x,0) \in \Lambda$ belongs to an atom $A \in \mathcal{A}_n$ that has an itinerary with $i_n = 1$, this itinerary is $(i_1, i_2, \ldots, *, 1)$ (where $*$ denotes 2 or 3). Then

$$(x,0) \in \{(x,y) \in X : y = 0\} \cap f_1\left(\overline{X_1} \cap \left(\overline{f_2(X_2)} \cup \overline{f_3(X_3)}\right)\right) = \{f_1(x_s, 0)\} \subset \mathcal{S}.$$

(ii) If $i_n = 5$, then either $i_{n-1} = 5$ of $i_{n-1} = 1$, because the only pieces X_i such that $\overline{f(X_i)} \cap X_5 \neq \emptyset$ are X_1 or X_5. So, if an atom $A \in \mathcal{A}_n$ has an itinerary such that $i_n = 5$, this itinerary either is $(i_1, i_2, \ldots, *, 1, 5, 5, 5, 5)$ (where the last $k \leqslant n$, $k \geqslant 1$ symbols are 5), or $(5, \ldots, 5)$, where all the symbols are 5. Then either $A = f_5^k(A')$, where A' is an atom in the case (i), or $A = f^n(X_5)$. Therefore, for all $(x,0) \in \Lambda$ such that $(x,0) \in A \in \mathcal{A}_n$ with $i_n = 5$, we have either $(x,0) = f_5^k \circ f_1(x_s, 0)$ for some $k \geqslant 1$, or $(x,0) \in \overline{f_5^k(X_5)}$ for all $k \geqslant 1$. We conclude that

$$(x,0) \in \bigcup_{k \geqslant 1} \{f_5^k \circ f_1(x_s,0)\} \subset \mathcal{S} \quad \text{or} \quad (x,0) = \bigcap_{k \geqslant 1} f_5^k(X_5) = \{(1,0)\} \subset \mathcal{S}.$$

(iii) If $i_n = 4$, then $i_{n-1} = 5$, because the only piece X_i such that $\overline{f(X_i)} \cap X_4 \neq \emptyset$ is X_5. So, if an atom A has itinerary such that $i_n = 4$, this itinerary is $(i_1, i_2, \ldots, 5, 4)$. Then $A = f_4(A')$ where A' is an atom of the case (ii). Therefore, for all $(x,0) \in \Lambda$ such that $(x,0) \in A \in \mathcal{A}_n$ with $i_n = 4$, we have

$$(x,0) \in \{(x,y) \in X : y = 0\} \ \cap \ f_4(\mathcal{S}) = \{f_4(1,0)\} = \{(x_e, 0)\} \subset \mathcal{K}.$$

(iv) If $i_n \in \{2,3\}$, then either $i_{n-1} = 4$ or $i_{n-1} \in \{2,3\}$, because the only pieces X_i such that $\overline{f(X_i)} \cap (X_2 \cup X_3) \neq \emptyset$ are X_5 or X_2 or X_3. So, if an atom A has itinerary such that $i_n \in \{2,3\}$, this itinerary is either $(i_1, \ldots, 4, 3)$ or $(i_1, \ldots, 4, 3, *, \ldots, *)$, where the last $k \geqslant 1$, $k \leqslant n-2$ symbols are $*$, or $(*, *, \ldots, *)$ (where the symbol $*$ denotes 2 or 3). Then, either $A = f_3(A' \cap \overline{X_3})$ or $A = g^k \circ f_3(A' \cap \overline{X_3}))$, for some $k \geqslant 1$, where A' is an atom of the case (iii), or $A \subset \overline{g^n(X_2 \cap X_3)}$, (recall from the fifth step of the construction of f that $f|_{X_i} = g|_{X_i}$ for $i = 2,3$). Therefore, for all $(x,0) \in \Lambda$ such that, for all $n \geqslant 1$, $(x,0) \in A \in \mathcal{A}_n$ with $i_n \in \{2,3\}$, we deduce:

$$(x,0) \in \{(x,y) \in X : y = 0\} \bigcap \left(\ \{f_3(x_e,0)\} \bigcup \left(\bigcup_{k \geqslant 1} \{g^k(f_3(x_e,0))\} \bigcup \left(\bigcap_{n \geqslant 1} \overline{g^n(X_2 \cap X_3)} \ \right) \ \right) \right) \subset \mathcal{K}.$$

The inclusion in the formula above is obtained as follows: First, $f(x_e, 0) = (h(x_e), 0) \in K \times \{0\}$, because $x_e \in K$ and after Theorem 3.9, $h(K) \subset K$. Second, after Formula (17) and by the fifth step in the construction of f we have $f_3(x_e, 0) = g(x_e, 0)$, so $g^k(f_3(x_e, 0)) = (F^{k-1}(x_e), 0)$. Using the same argument as above $h^{k-1}(x_e) \in K$ for all $n \geqslant 0$. Then $g^k(f_3(x_e, 0)) \in K \times \{0\}$ for all $k \geqslant 1$. Third, after Formula (17),

$$\{(x, y) \in X : y = 0\} \bigcap \{ \bigcap_{n \geqslant 1} \overline{g^n(X_2 \cap X_3)} \} = \{ \bigcap_{n \geqslant 1} h^n([0, 1]) \} \times \{0\} = K \times \{0\},$$

because K is the attractor of h, as stated in Theorem 3.9.

As any point $(x, y) \in \Lambda$ is necessarily included in some atom A of generation n, for all $n \geqslant 1$, it belongs to one of the above four cases. Therefore $(x, y) \in \mathcal{K} \cup \mathcal{S}$. □

Lemma 3.14.

$$\textit{For some } (x_0, y_0) \in X_5 \cap \tilde{X} \textit{ such that } y_0 > 0 : \ \omega(x_0, y_0) \supset \mathcal{K} \cup \mathcal{S}. \tag{24}$$

Proof. We shall prove that $\mathcal{K} \subset \omega(x_0, y_0)$ and that $\mathcal{S} \subset \omega(x_0, y_0)$.

Consider the open set $\mathcal{D}$ constructed in Lemma 3.11. Choose some $(x_0, y_0) \in \tilde{X} \cap \mathcal{D}$. Such a point exists because $\tilde{X}$ is dense in X. (See Theorem 2.8.) Denote

$$(x_n, y_n) := f^n(x_0, y_0) \, \forall n \geqslant 0 \quad \text{and} \quad \mathcal{O} := \{(x_n, y_n)\}_{n \geqslant 0}.$$

It is easy to check, taking into account the formula of each of the homeomorphisms f_i (defined in the fifth step of the construction of f) that:

$$y_n > y_{n+1} > 0 \ \ \forall \ n \geqslant 0 \quad \text{and} \quad \lim_{n \to +\infty} y_n = 0. \tag{25}$$

Applying Lemma 3.11 to the point $(x_0, y_0) \in \mathcal{D} \cap \tilde{X}$, there exist some n such that $f^n(x_0, y_0) \in X_1$. So, the orbit $\mathcal{O}$ visits the piece X_1 at least once, after it visits the open set $\mathcal{D}$. Since $f(X_1) \subset X_5$, the orbit $\mathcal{O}$ visits the piece X_5, at least once, after it visits the piece X_1. Since $f|_{X_5} = f_5$ is a contracting homothecy with center at $(1, 0)$, the half line $\{(x, y) \in X : x \geqslant 1, y = 0\}$ containing any point $(x_n, y_n) \in X_5$ also contains the iterates $f^k(x_n, y_n)$ for all $k \geqslant 1$ such that $f^{k-1}(x_n, y_n) \in X_5$. But, because that parabola has an horizontal tangent direction at the point $(1, 0)$, that line intersects the parabolic arc $\Delta_{45} = \partial X_4 \cap \partial X_5$ (see Figure 4). We deduce that if $(x_n, y_n) \in X_5 \cap \tilde{X}$ satisfies $y_n > 0$, then there exists $k \geqslant 1$ such that $(x_{n+k}, y_{n+k}) \in X_4$. Since $y_n > 0$ for all the points of the orbit $\mathcal{O}$ that we are studying, it visits the piece X_4 at least once after visiting the piece X_5. Since by definition $f(X_4) \subset \mathcal{D}$, the orbit $\mathcal{O}$ visits $\mathcal{D}$ at least once after it visits the piece X_4. Applying all the arguments again to the point $(x_n, y_n) \in \mathcal{D} \cap \tilde{X}$, so obtained for some $n > 1$ (recall that $f(\tilde{X}) \subset \tilde{X}$) we obtain that for all $i \in \{1, 4, 5\}$ there exist infinitely many values of n such that $(x_n, y_n) \in X_i$, infinitely many values of n such that $(x_n, y_n) \in \mathcal{D} \subset X_3$, and those sets are visited by the orbit $\mathcal{O}$ in the following order: $(\mathcal{D}, X_1, X_5, X_4, \mathcal{D}, X_1, X_5, X_4, \ldots)$, but not necessarily in consecutive iterates (i.e. the time during which the orbit stays in some of the pieces is not necessarily 1).

Define the sequences of *return times* $n_j^{(\cdot)}$ to each of the sets X_5 and $\mathcal{D}$, and the sequences of the *staying times* $m_j^{(\cdot)}$ in the sets X_5 and $X_2 \cap X_3$:

$$n_1^{(5)} := \min\{n \geqslant 1 : (x_n, y_n) \in X_5\}, \quad m_1^{(5)} := \min\{m \geqslant 1 : f^m(x_{n_1^{(5)}}, y_{n_1^{(5)}}) \notin X_5\}.$$

By induction, for all $j \geqslant 1$, we define:

$$n_{j+1}^{(5)} := \min\{n \geqslant n_j^{(5)} + m_j^{(5)} : (x_n, y_n) \in X_5\}, \quad m_{j+1}^{(5)} := \min\{m \geqslant 1 : f^m(x_{n_{j+1}^{(5)}}, y_{n_{j+1}^{(5)}}) \notin X_5\}.$$

Analogously:

$$n_1^{(\mathcal{D})} := \min\{n \geqslant 0 : (x_n, y_n) \in \mathcal{D}\} = 0, \quad m_1^{(2,3)} := \min\{m \geqslant 1 : f^m(x_{n_1^{(\mathcal{D})}}, y_{n_1^{(\mathcal{D})}}) \notin X_2 \cup X_3\}.$$

By induction, for all $j \geqslant 1$, we define:

$$n_{j+1}^{(\mathcal{D})} := \min\{n \geqslant n_j^{(\mathcal{D})} + m_j^{(2,3)} : (x_n, y_n) \in \mathcal{D}\}, \quad m_{j+1}^{(2,3)} := \min\{m \geqslant 1 : f^m(x_{n_{j+1}^{(\mathcal{D})}}, y_{n_{j+1}^{(\mathcal{D})}}) \notin X_2 \cup X_3\}.$$

For each $j \geqslant 1$ consider the *escape point*

$$(\widehat{x}_j, \widehat{y}_j) := f^{m_j^{(2,3)}}(\tilde{x}_j, \tilde{y}_j) \in X_1, \tag{26}$$

where $(\tilde{x}_j, \tilde{y}_j)$ is some *return point*. By definition of the staying time $m_j^{(2,3)}$ in the open set $X_2 \cap X_3$, and since $f|_{X_2 \cup X_3} = g$, the escape point can be computed using the Formula (17). Then

$$\widehat{y} = \alpha^{m_j^{(2,3)}}\, \tilde{y}_j \leqslant \alpha\, \tilde{y}_j$$

Because of the previously determined order, the consequent $(i+1)$–th return point $(\tilde{x}_{j+1}, \tilde{y}_{j+1})$ of the orbit $\mathcal{O}$ to the open set $\mathcal{D}$ will not appear before the orbit had escaped from $X_2 \cap X_3$ and visited X_1 at the point $(\widehat{x}_j, \widehat{y}_j)$. From Inequality (25) we have $y_{n+1} < y_n$ for all $n \geqslant 0$. So:

$$\tilde{y}_{j+1} < \widehat{y}_j \leqslant \alpha\, \tilde{y}_j \quad \forall\ j \geqslant 1.$$

Therefore, by induction on $j \geqslant 1$ we obtain $\tilde{y}_j \leqslant \alpha^{j-1} \tilde{y}_1 = y_0$ for all $j \geqslant 1$. Taking logarithm and computing the limit, this last inequality implies the so called *sequence of return points* to $\mathcal{D}$ (i.e. the orbit by the return map to $\mathcal{D}$) defined by:

$$\{(\tilde{x}_j, \tilde{y}_j) := f^{n_j^{(\mathcal{D})}}(x_0, y_0)\}_{j \geqslant 1}$$

satisfies:

$$\lim_{j \to +\infty} \frac{\log \tilde{y}_j}{j} \leqslant \log \alpha.$$

After part c) of Lemma 3.11, it follows that $\lim_{j \to +\infty} m_j^{(2,3)} = +\infty$. Indeed, since the staying time $m_j^{(2,3)}$ of the orbit $\mathcal{O}$ in the pieces $X_2 \cup X_3$ is, by definition, the minimum integer number m such that the g^m–iterate of the return point $(\tilde{x}_j, \tilde{y}_j) \in \mathcal{D}$ drops in the piece X_1.

Consider the return point $(\tilde{x}_j, \tilde{y}_j)$ for some $j \geqslant 1$. Fix any $p = (x_p, 0) \in K$, fix a positive and arbitrarily small value of $\epsilon < 1$, and consider the open ball $B_p(\epsilon/2)$. From Equality (21), there exists a natural number k such

$$k > \frac{\log 4 - \log \epsilon}{-\log \alpha},$$

and such that the iterate

$$f^k(x_e, 0) = g^k(x_e, 0) \in B_p(\epsilon/2).$$

Since the sequence $m_j^{(2,3)}$, formed by the staying times in $X_2 \cup X_3$ of the orbit $\mathcal{O}$, is divergent, there exists some $j_0 \geqslant 1$ such that $m_{j_0}^{(2,3)} > k$. Recalling that $(\tilde{x}_j, \tilde{y}_j)$ is the j–th return point of the orbit $\mathcal{O}$ to $\mathcal{D} \subset X_3$, and considering the above order of visits, we can write:

$$(\tilde{x}_j, \tilde{y}_j) \in \mathcal{D}, \ \ 1 \leqslant k < m_{j_0}^{(2,3)} = \min\{m : f^m(\tilde{x}_j, \tilde{y}_j) \in X_1\}.$$

Applying the statement b) of Lemma 3.11 we obtain:

$$d(\,(f^k(\tilde{x}, \tilde{y}),\ f^k(x_e, 0)\,) \leqslant d(\,(f^k(\tilde{x}, \tilde{y}),\ f^k(x_e, \tilde{y})\,) + d(\,(f^k(x_e, \tilde{y}),\ f^k(x_e, 0)\,) \leqslant$$

$$\leqslant \alpha^k + \alpha^k\, \tilde{y} \leqslant 2\,\alpha^k = \exp(\log 2 + k \log \alpha) < \exp(\log 2 + \log \epsilon - \log 4) = \frac{\epsilon}{2}.$$

Therefore:

$$d(f^k(\tilde{x}, \tilde{y}), (x_p, 0)) \leqslant d(f^k(\tilde{x}, \tilde{y}), f^k(x_e, 0)) + d(f^k(x_e, 0), (x_p, 0)) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Thus, for all $0 < \epsilon < 1$ the orbit $\mathcal{O}$ of the point (x_0, y_0), has an iterate at distance smaller than ϵ of the point $p = (x_p, 0) \in K$, that is to say $(x_p, 0) \in \omega(x_0, y_0)$. As the argument above works for any point $p \in \mathcal{K}$ we deduce that

$$\mathcal{K} \subset \omega(x_0, y_0).$$

Consider the sequence of escaping points $(\widehat{x}_j, \widehat{y}_j)$ of the orbit $\mathcal{O}$, from the pieces $X_2 \cup X_3$. After determined the visit order, they are the return points to the piece X_1. From the definition of return times and staying times in the part 3. of this proof, we have:

$$(\widehat{x}_j, \widehat{y}_j) = f^{m_j^{(2,3)}}(\tilde{x}_j, \tilde{y}_j) \in X_1, \qquad (\tilde{x}_j, \tilde{y}_j) \in \mathcal{D}, \qquad m_j^{(2,3)} = \min\{m \geqslant 1 : f^m(\tilde{x}_j, \tilde{y}_j) \in X_1\}.$$

The divergence of staying times allows us to use the part d) of Lemma 3.11 to deduce that

$$\lim_{j \to +\infty} (\widehat{x}_j, \widehat{y}_j) = (x_s, 0).$$

As $(\widehat{x}_j, \widehat{y}_j) \in X_1$ and f_1 is continuous, we obtain:

$$\lim_{j \to +\infty} f(\widehat{x}_j, \widehat{y}_j) = \lim_{j \to +\infty} f_1(\widehat{x}_j, \widehat{y}_j) = f_1(x_s, 0).$$

Therefore, the orbit $\mathcal{O}$ has infinitely many points of the form $f(\widehat{x}_j, \widehat{y}_j)$ in any arbitrarily small neighborhood of $f_1(x_s, 0)$. In other words

$$f_1(x_s, 0) \in \omega(x_0, y_0).$$

Since the orbit of $f_1(x_s, 0)$ by f_5 remains in the open contractive piece X_5, we have that f^k is continuous in the point $f_1(x_s, 0)$ for all $k \geqslant 1$. Therefore, if a subsequence of the orbit $\mathcal{O}$ converges to $f_1(x_s, 0)$, the f^k–iterates of that subsequence converges to $f^k \circ f_1(x_s, 0) = f_5^k \circ f_1(x_s, 0)$ for any fixed $k \geqslant 1$. We conclude that

$$\{f_5^k \circ f_1(x_s, 0)\}_{k \geqslant 0} \subset \omega(x_0, y_0).$$

Finally as $\omega(x_0, y_0)$ is closed, applying Equality (22) which states that the closure of the set $\{f_5^k \circ f_1(x_s, 0)\}_{k \geqslant 0}$ is $\mathcal{S}$, we conclude that

$$\mathcal{S} \subset \omega(x_0, y_0).$$

□

Proof of Proposition 3.12. From Lemma (3.14), Theorem 2.9 and Lemma (3.13) we have the following chain of inclusions:

$$\mathcal{K} \cup \mathcal{S} \ \subset \ \omega(x_0, y_0) \ \subset \ \bigcup_{(x,y) \in \tilde{X}} \omega(x, y) = L \ \subset \ \Omega \ \subset \ \Lambda \ \subset \ \mathcal{K} \cup \mathcal{S}.$$

□

4 Conclusion and Further Questions

In this paper we have adapted several definitions and obtained general results (Theorems 2.9, 2.10 and 2.12) to characterize the attractors of piecewise contracting maps acting on any compact and locally connected metric space. The class of systems studied here can fit a lot of concrete models, in particular those considered so far since it includes any piecewise contracting C^1 map on any finite dimensional manifold, and therefore any piecewise affine map on a (compact) Euclidean space. But actually, for our results to hold, the map only needs to be piecewise C^0 contracting (and can include directions of infinite contraction when differentiable), and the phase space can be for instance a piece of any topological vector space or any infinite dimensional as well as dimensionless manifold. The contraction has to take place on any finite number of open pieces, but the frontiers between those pieces are not necessarily piecewise C^1, nor necessarily have zero Lebesgue measure (whenever the latter is defined).

The chief property that has been investigated in the previous literature is the generic asymptotic periodicity of piecewise contracting maps. The proofs consist in showing the genericity of the property $d(\Lambda, \Delta) > 0$, or a similar one, and showing that it implies periodicity. Both parts of the proof have been done so far in specific contexts. The contribution of Theorem 2.10 is to generalize the proof of the periodicity result to a wider class of maps that encompass all the previous models and others where classical arguments cannot be applied. This theorem gives a practical criterion to verify periodicity for a great number of applied models. However, the question of whether the periodic behavior is C^∞, or at least C^1 generic (with respect to the differentiability of the pieces of Δ) remains open.

When $d(\Lambda, \Delta) = 0$, we found attractors that are finite, countably infinite, or uncountable but totally disconnected (cf. Propositions 3.1, 3.3, 3.5, 3.7, 3.9, 3.12). Thus we obsolete the dichotomy observed so far according to which the asymptotic dynamics is either supported by a finite set of periodic orbits or by a set which essentially consists of a Cantor set. We achieved to show the diversity of possible attractors and unexpected recurrence properties just working in compact rectangles of $\mathbb{R}^2$, with piecewise analytic frontiers (parabolic arcs and line segments) except in Example 3.6, and which is more surprising, with maps that are simply piecewise affine. Therefore, we cannot exclude the possibility to find more complex attractors, having for example connected components or infinitely many transitive components, considering wilder maps (violating necessarily all the hypothesis of Theorem 2.12 for the former example). On the other hand, in all known examples, including ours, either the attractor is a finite union of periodic orbits or it is not persistent under arbitrarily small perturbations of the map. Therefore, the presented results deal, as far as we can tell, with the class of bifurcating dynamics. A better understanding of the case $d(\Lambda, \Delta) = 0$ is part of a complete characterization of bifurcations, in particular border–collision bifurcations, in families of piecewise contracting maps when unfolded with some parameters.

The present paper, specifically its section 3, shows that the property of the dynamics being contracting everywhere except perhaps in a topologically negligible set (see Theorem 2.8) determines only a few proper characteristics of the attractor. Even more, we have shown that the additional condition $d(\Lambda, \Delta) > 0$ is in general only sufficient for the finiteness of Λ (see Proposition 3.1). In order to obtain more discernible dynamical properties, the necessity to determine subclasses by imposing further, adequate restrictions on this general class of maps becomes evident. Considering restrictions is essentially what has been done in the literature so far, without being aware of the diversity of behaviors this class of maps offers though. As mentioned above, several open questions naturally appear now in relation to the bifurcations and complexity in this class of maps. Answering those questions may in particular show their potential use to model complex systems in engineering as well as in the natural and social sciences.

Acknowledgements. EC was partially financed by CSIC of Universidad de la Repblica and ANII of Uruguay. PG has been supported by grant of Chile FONDECYT 1100764. AM was partially financed by the scientific cooperation between the IVIC and the UdelaR. EU gratefully acknowledges SEP–Mexico. During the development of this work, EC, AM and EU were invited at the Universidad de Vaparaíso and benefited from the financial support of FONDECYT 1100764. AM and EU were also invited at the Universidad de la República.

5 Appendix

The purpose of this appendix is to prove that there exists piecewise contracting maps, according to Definition 2.1, exhibiting atoms of arbitrarily large generation that have (infinitely) many connected components. The following example shows that a connection and, more importantly and not considered in [4], the finiteness property of the atoms cannot be argued under our general hypothesis and therefore justifies the necessity of the proofs of Theorems 2.9, 2.10 and 2.12.

Consider the compact square $X = [-1, 1] \times [0, 1/e^{2\pi}] \subset \mathbb{R}^2$. Construct, as in Figure 5, the following two open pieces:

$$X_1 = \{(x, y) \in X : y = 0, x > 0\} \ \cup \ \{(x, y) \in X : y > 0, x > y \sin(-\log y)\}$$

$$X_2 = \{(x, y) \in X : y = 0, x < 0\} \ \cup \ \{(x, y) \in X : x < y \sin(-\log y)\}.$$

The set $\Delta = \partial X_1 \cap \partial X_2$ is the continuous compact arc:

$$\Delta = \{(0, 0)\} \ \cup \ \{x = y \sin(-\log |y|)\,,\ 0 < y \leqslant 1/e^{2\pi}\},$$

which is the graph of a continuous real function $x(y)$ defined in a compact interval of real values of y. (Notice that $\lim_{y \to 0} x(y) = 0$.)

Define the contracting maps:

$$f_1(x, y) = \frac{1}{e^{\pi}}(x, y), \quad \text{and} \quad f_2(x, y) = \frac{-1}{2} + \frac{1}{2}\left(x + \frac{1}{2}, y\right).$$



Figure 5: For all generations greater than 2, each one of the atoms $A_{12} = \overline{f(\overline{f(X_1)} \cap X_2)}$, $A_{122} = f(A_{12}), \ldots A_{122\ldots 2} = f^n(A_{12})$ has infinitely many connected components. In this example we have $\Lambda = \Omega = L = \{(-1/2, 0),\ (0, 0)\}$.

Construct any map $f : X \mapsto X$ such that $f|_{X_1} = f_1|_{X_1}$ and $f|_{X_2} = f_2|_{X_2}$, as shown in Figure 5. Then, f is a piecewise (affine) contracting map according to our Definition 2.1.

Proposition 5.1. *The piecewise contractive map f constructed above exhibits, for all $n \geqslant 2$, an atom A_n of generation n which is disconnected and such that the connected components of A_n are infinitely many.*

Proof. By construction, f_2 is an homeomorphism, and besides $\overline{f_2(X_2)} = f_2(\overline{X_2}) \subset X_2$. Therefore, if an atom A of generation n_0 has infinitely many connected components and is contained in $\overline{X_2}$, then for all $n > n_0$ the atoms $\overline{f^{n-n_0}(A)} = f_2^{n-n_0}(A)$ of generation n are homeomorphic images of A contained in X_2. Thus, they also have infinitely many connected components. So, to end the proof of Proposition 5.1, it is enough to check that the atom A_{12} of generation 2 satisfies $A_{1,2} \subset \overline{X_2}$, and has infinitely many connected components.

Taking into account Definition 2.4:

$$A_1 = \overline{f_1(X_1)} = \{(x', y') \in X : x' \geqslant y' \sin(\pi - \log y')\},$$

$$A_{12} = \overline{f_2(A_1 \cap X_2)} = f_2(A_1 \cap \overline{X_2}) = \{0, 0\} \cup f_2(B), \text{ where}$$

$$B = \{(x, y) \in \mathbb{R}^2 : 0 < y \leqslant 1/e^{2\pi},\ \ y \sin(\pi - \log y) \ \leqslant x\ \leqslant y \sin(-\log y)\}.$$

Observe that the interval $(0, e^{-2\pi}]$ can be written as the following disjoint union:

$$(0, e^{-2pi}] = \bigcup_{j=1}^{+\infty} \left(\{1/e^{(2j+1)\pi} \leqslant y \leqslant 1/e^{2j\pi}\} \cup \{1/e^{(2j+2)\pi} < y < 1/e^{(2j+1)\pi}\} \right),$$

and observe that for all natural number $j \geqslant 1$ the following set is empty:

$$C_j = \Big\{ (x, y) \in X : (1/e^{(2j+2)\pi}) < y < (1/e^{(2j+1)\pi}), \text{ and } \sin(\pi - \log y) \leqslant x \leqslant y \sin(-\log y) \Big\} = \emptyset,$$

because $\sin(\pi + \alpha) > 0 > \sin(\alpha)$ if $(2j + 1)\pi < \alpha = -\log y < (2j + 2)\pi$.

Therefore we deduce that

$$A_{12} = \{f_2(0,0)\} \bigcup \Bigg(\bigcup_{j=1}^{\infty} f_2(B_j) \Bigg), \text{ where} \tag{27}$$

$$B_j = \Big\{ (x,y) \in X : (1/e^{(2j+1)\pi}) \leqslant y \leqslant (1/e^{2j\pi}), \text{ and } \sin(\pi - \log y) \leqslant x \leqslant y \sin(-\log y) \Big\}.$$

Notice that the family $\{B_j\}_{j \geqslant 1}$ is composed of infinitely many connected compact sets contained in $\overline{X_2}$ that are pairwise disjoint (the same values of y cannot be shared by the points of B_j and B_i if $j \neq i$). Since f_2 is an homeomorphism and $f_2(\overline{X_2}) \subset X_2$, we deduce that the family $\{f_2(B_j)\}_{j \geqslant 1}$ is also composed by infinitely many pairwise disjoint connected compact sets contained in X_2. So the Equality (27) implies that the atom A_{12} is disconnected, has infinitely many connected components, and is contained in X_2 as wanted. □

References

[1] Avrutin V and Schanz M 2008 On the fully developed bandcount adding scenario *Nonlinearity* **21** (1) 1077–1103

[2] Blokhina E and Feely O 2009 A kicked oscillator as a model of a pulsed MEMS system *International Journal of Bifurcation and Chaos* **19** (1) 187–202

[3] Bremont J 2006 Dynamics of injective quasi-contractions *Ergodic Theory and Dynamical Systems* **26** 19–44

[4] Bruin H and Deane J 2009 Piecewise contractions are asymptotically periodic *Proceedings of the American Mathematical Society* **137** (4) 1389–1395

[5] Buzzi J 2001 Piecewise isometries have zero topological entropy *Ergodic Theory and Dynamical Systems* **21** 1371–1377

[6] Casey R, de Jong H and Gouzé J-L 2006 Piecewise–linear models of genetic regulatory networks: equilibria and their stability *Journal of Mathematical Biology* **52** (1) 27–56

[7] Catsigeras E 2010 Chaos and Stability in a Model of Inhibitory Neuronal Network *International Journal of Bifurcation and Chaos* **20** (2) 349–360

[8] Catsigeras E and Budelli R 2011 Topological dynamics of generic piecewise contractive maps in n dimensions *International Journal of Pure and Applied Math.* In press. arXiv [math.DS] 0805.2695v2.

[9] Catsigeras E and Guiraud P 2010 Integrate and Fire Neural Networks, Piecewise Contractive Maps and Limit Cycles arXiv:1011.1525v1

[10] Cessac B 2008 A discrete time neural network model with spiking neurons. Rigorous results on the spontaneous dynamics *Journal of Mathematical Biology* **56** (3) 311–345

[11] Coutinho R, Fernandez B, Lima R and Meyroneinc A 2006 Discrete time piecewise affine models of genetic regulatory networks *Journal of Mathematical Biology* **52** (4) 524–570

[12] Cros A, Morante A and Ugalde E 2008 Regulatory dynamics on random networks: asymptotic periodicity and modularity *Nonlinearity* **21** (3) 537–556

[13] Edwards R, van den Driessche P and Wang L 2007 Periodicity in piecewise–linear switching networks with delay *Journal of Mathematical Biology* **55** (2) 271–298

[14] Feely O and Chua L O 1991 The effect of integrator leak in modulation *IEEE Transactions on Circuits and Systems* **38** (11) 1293–1305

[15] Galatolo S 2003 Complexity initial condition sensitivity dimension and weak chaos in dynamical systems *Nonlinearity* **16** (4) 1219–1239

[16] Galatolo S 2003 Global and local complexity in weakly chaotic dynamical systems *Discrete and Continuous Dynamical Systems* **9** (6) 1607–1624

[17] Gambaudo J–M and Tresser Ch 1988 On the dynamics of quasi-contractions *Bulletin of the Brazilian Mathematical Society* **19** (1) 61–114

[18] Goetz A and Poggiaspalla G 2004 Rotations by $\pi/7$ *Nonlinearity* **17** (5) 1787–1802

[19] Katok A and Hasselblatt B 1995 Introduction to the modern theory of dynamical systems *Encyclopedia of Mathematics and its Applications* vol 54 (Cambridge University Press)

[20] Kechris A S 1995 Classical Descriptive Set Theory *Graduate Texts in Mathematics* vol 156 (Springer)

[21] Keener J P 1980 Chaotic behavior in piecewise continuous difference equations *Transactions of the American Mathematical Society* **261** (2) 589–604

[22] Keller G and Liverani C 2006 Uniqueness of the SRB measure for piecewise expanding weakly coupled map lattices in any dimension *Communications in Mathematical Physics* **262** (1) 33–50

[23] Lima R and Ugalde E 2006 Dynamical complexity of discrete–time regulatory networks *Nonlinearity* **19** (1) 237–259

[24] Lowenstein J H, Poggiaspalla G and Vivaldi F 2005 Sticky orbits in a kicked–oscillator model *Dynamical Systems* **20** (4) 413–451

[25] Mirollo R E and Strogatz S H 1990 Synchronization of pulse-coupled biological oscillators *SIAM Journal of Applied Mathematics* **50** (6) 1645–1662

[26] Mosekilde E and Zhusubalyev Zh 2003 Bifurcations and Chaos in Piecewise–Smooth Dynamical Systems *World Scientific Series on Nonlinear Science* Series A vol 44 (World Scientific)

[27] Palis J and Takens F 1993 Hyperbolicity and Sensitive Chaotic Dynamics of Homoclinic Bifurcations *Cambridge Studies in Advanced Mathematics* No 35 (Cambridge University Press)

[28] Tramontana F, Gardini L and Ferri P 2010 The dynamics of the NAIRU model with two switching regimes *Journal of Economic Dynamics and Control* **34** (4) 681–695

[29] Kruglikov B and Rypdal M 2006 A piecewise affine contracting map with positive entropy *Discrete and Continuous Dynamical Systems* **16** (2) 393–394